

AM- 80- 129

American Mineralogist, Volume 65, pages 557-562, 1980

Fluorine distribution in the hydrous silicate minerals of the Franklin Marble, Orange County, New York

LANCE E. KEARNS, LUCILLE E. KITE¹

Department of Geology, James Madison University
Harrisonburg, Virginia 22807

PETER B. LEAVENS

Department of Geology, University of Delaware
Newark, Delaware 19711

AND JOSEPH A. NELEN

Department of Mineral Sciences, Smithsonian Institution
Washington, D. C. 20560

Abstract

Fifty specimens of the hydrous silicates norbergite, chondrodite, phlogopite, tremolite, edenite, and pargasite from 34 localities in the Franklin Marble of Orange County, New York, were analyzed quantitatively, and their F/(F + Cl + OH) ratios (X_F) determined. The average F/(F + Cl + OH) ratio (\bar{X}_F) determined for each species follows the sequence: Trem < \bar{X}_F Phlog < \bar{X}_F Amph < \bar{X}_F Chond < \bar{X}_F Norb where \bar{X}_F Amph is the average X_F for amphibole species other than tremolite. The fluorine contents of the hydrous silicate minerals formed within the Franklin Marble are high by comparison with analyses of the same species from other localities. The Franklin Marble is a high-temperature ($836 \pm 40^\circ\text{C}$), high-pressure (4–7 kbar) metacarbonate. We suggest that the high fluorine content of the tremolite samples has increased the upper thermal stability limit for this species, thus preserving the assemblage tremolite + quartz + calcite under granulite facies conditions. There is an inverse relationship between the F/(F + Cl + OH) ratio and titanium content for chondrodite. This relationship has been noted by earlier investigators for other members of the humite group. The inverse relationship is not evident for Ti values below 0.02 atoms per formula unit.

Introduction

The Franklin Marble is a late Precambrian (Grenvillian age) metacarbonate which crops out in a narrow belt in Sussex County, New Jersey and Orange County, New York, as part of the Hudson Highlands. Hague *et al.* (1956) and Offield (1967) have mapped portions of the Highlands and determined the conditions of metamorphism to be upper amphibolite to granulite facies. However, a number of hydrous silicate species, including tremolite and other amphiboles, norbergite, and chondrodite, occur

widely in the marble. The occurrence of tremolite in particular is not compatible with the postulated metamorphic conditions.

Earlier work on the humites of the Franklin Marble by Jones *et al.* (1969) suggests that the hydrous silicates are rich in fluorine. We undertook to determine the amount and distribution of fluorine in the hydrous silicates of the Franklin Marble in Orange County, New York, and to study the relationship of fluorine content to the stability of hydrous silicates under conditions of high-grade metamorphism. We conclude that high fluorine contents stabilize the hydrous silicates under these conditions and are responsible for their persistence in this granulite facies metacarbonate.

¹ Present address: Department of Geoscience, North Carolina State University, Raleigh, North Carolina 27607.

ANALYSES OF HUMITES, AMPHIBOLES AND PHLOGOPITE
FROM THE FRANKLIN MARBLE, ORANGE COUNTY, NEW YORK

Lance E. Kearns
Department of Geology, James Madison University
Harrisonburg, Virginia 22807

Lucille E. Kite
Department of Geology, James Madison University
Harrisonburg, Virginia 22807

Peter B. Leavens
Department of Geology, University of Delaware
Newark, Delaware 19711

Joseph A. Nelen
Department of Mineral Science, Smithsonian Institution
Washington, D. C. 20560

Analytical Results

Humites

The humite specimens were all analyzed by electron microprobe which yielded weight oxide percents for the cations (Si, Al, Ti, Mg, Fe, Mn) and for F and Cl.

Calculation of the chemical formula from the mineral analysis was done according to Deer, Howie and Zussman (1966), and based on six oxygen equivalents for the norbergite and ten oxygen equivalents for chondrodite. The atomic proportion of OH was determined by the formula (Jones, Ribbe and Gibbs, 1969):

$$\text{OH} = (2/(2n + 1)) M_{\text{Ti}} - 2\text{Ti} - \text{F}$$

where $n = 1$ for norbergite, $n = 2$ for chondrodite, and M_{Ti} represents the sum of the atomic proportions of Fe, Mn, Mg, and Ti.

From this atomic proportion of OH the weight of H_2O^+ was calculated.

The oxide percent total in some cases is somewhat lower than one would expect for a complete analysis and is perhaps due to the presence of small amounts of Na or K which were not analyzed for in this study. (See Tables 1a and 1b for humite analyses.)

Table 1a.
Index of Specimens for Humite Analyses.

| Analysis | Sample #* | Mineral Species |
|----------|-----------|-----------------|
| 1 | H-15 | Norbergite |
| 2 | H-56 | Norbergite |
| 3 | H-64 | Norbergite |
| 4 | H-55A | Norbergite |
| 5 | H-132 | Chondrodite |
| 6 | H-15A | Chondrodite |
| 7 | H-200-44 | Chondrodite |
| 8 | H-157 | Chondrodite |
| 9 | H-20 | Chondrodite |
| 10 | H-94 | Chondrodite |
| 11 | H-143 | Chondrodite |
| 12 | H-90 | Chondrodite |

*Indicates mineral locality (See Figure 1)

Table 1b.

Analyses of Humites

| | #1 | #2 | #3 | #4 | #5 | #6 |
|--------------------------------|-----------------|-----------------|-----------------|-----------------|-----------------|-------------|
| SiO ₂ | 29.17 | 29.21 | 28.25 | 28.92 | 33.63 | 33.74 |
| Al ₂ O ₃ | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| TiO ₂ | 0.47 | 0.23 | 0.11 | 0.28 | 0.36 | 0.30 |
| MgO | 58.02 | 58.51 | 58.61 | 57.93 | 51.09 | 53.44 |
| FeO | 0.57 | 0.21 | 0.41 | 0.40 | 5.19 | 3.07 |
| MnO | 0.05 | 0.06 | 0.03 | 0.02 | 0.17 | 0.17 |
| H ₂ O ⁺ | 2.72 | 2.71 | 2.81 | 2.74 | 1.64 | 1.77 |
| F | 12.47 | 12.55 | 12.48 | 12.33 | 6.59 | 6.57 |
| Cl | <u><0.01</u> | <u><0.13</u> | <u><0.01</u> | <u><0.01</u> | <u><0.01</u> | <u>0.01</u> |
| Total | 103.47 | 103.61 | 102.70 | 102.62 | 98.67 | 99.06 |
| F+Cl=0 | <u>5.25</u> | <u>5.31</u> | <u>5.25</u> | <u>5.19</u> | <u>2.77</u> | <u>2.77</u> |
| Corrected Total | 98.22 | 98.29 | 97.44 | 97.43 | 95.90 | 96.29 |

Ions per formula unit based on 6(O, OH, F) for analyses

1-4, and 10(O, OH, F) for analyses 5,6

| | | | | | | |
|--------------------|------|------|------|------|------|------|
| Si | 1.00 | 1.00 | 0.98 | 1.01 | 2.05 | 2.02 |
| Al | -- | -- | -- | -- | -- | -- |
| Ti | 0.01 | -- | 0.01 | 0.01 | 0.02 | 0.01 |
| Mg | 2.97 | 2.99 | 3.02 | 3.02 | 4.63 | 4.78 |
| Fe | 0.02 | 0.01 | 0.01 | 0.01 | 0.26 | 0.15 |
| Mn | -- | -- | -- | -- | 0.01 | 0.01 |
| OH | 0.62 | 0.62 | 0.65 | 0.64 | 0.67 | 0.71 |
| F | 1.35 | 1.36 | 1.36 | 1.37 | 1.27 | 1.25 |
| Cl | -- | 0.01 | -- | -- | -- | -- |
| Ratio F F+Cl+OH | .68 | .68 | .68 | .63 | .65 | .64 |

| | #7 | #8 | #9 | #10 | #11 | #12 |
|--------------------------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| SiO ₂ | 33.39 | 34.17 | 33.82 | 33.23 | 33.37 | 33.34 |
| Al ₂ O ₃ | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| TiO ₂ | 1.08 | 0.15 | 1.18 | 0.73 | 0.44 | 0.27 |
| MgO | 51.61 | 55.61 | 54.92 | 53.55 | 56.20 | 57.19 |
| FeO | 4.78 | 1.84 | 1.97 | 3.78 | 1.75 | 0.91 |
| MnO | 0.15 | 0.05 | 0.06 | 0.21 | 0.10 | 0.05 |
| H ₂ O ^F | 1.88 | 2.19 | 2.00 | 1.86 | 1.89 | 2.06 |
| F | 5.86 | 6.00 | 5.89 | 6.32 | 6.63 | 6.43 |
| Cl | <u><0.01</u> | <u><0.01</u> | <u><0.01</u> | <u><0.01</u> | <u><0.01</u> | <u><0.01</u> |
| Total | 98.75 | 100.01 | 99.84 | 99.68 | 100.88 | 100.25 |
| F+Cl=0 | <u>2.47</u> | <u>2.53</u> | <u>2.48</u> | <u>2.66</u> | <u>2.79</u> | <u>2.71</u> |
| Corrected Total | 96.28 | 97.48 | 97.36 | 97.02 | 98.09 | 97.54 |

Ions per formula unit based on 10(O, OH, F)

| | | | | | | |
|---------------------------|------|------|------|------|------|------|
| Si | 2.02 | 2.01 | 2.00 | 1.99 | 1.99 | 1.95 |
| Al | -- | -- | -- | -- | -- | -- |
| Ti | 0.05 | 0.01 | 0.05 | 0.03 | 0.02 | 0.01 |
| Mg | 4.66 | 4.88 | 4.84 | 4.78 | 4.91 | 5.01 |
| Fe | 0.24 | 0.09 | 0.10 | 0.19 | 0.09 | 0.04 |
| Mn | 0.01 | -- | -- | 0.01 | -- | -- |
| OH | 0.76 | 0.85 | 0.79 | 0.74 | 0.74 | 0.81 |
| F | 1.12 | 1.12 | 1.10 | 1.20 | 1.23 | 1.19 |
| Cl | -- | -- | -- | -- | -- | -- |
| Ratio $\frac{F}{F+Cl+OH}$ | .51 | .56 | .58 | .62 | .62 | .60 |

Amphiboles

The complete analyses of the amphibole specimens were obtained in three ways. Seven were analyzed by electron microprobe (denoted by * under Analytical Methods in Table 2a) which yielded weight oxide percents for the cations (Si, Al, Ti, Fe, Mg, Mn, Ca, Na, K) and for the anions F and Cl. Four were analyzed by electron microprobe to obtain weight oxide percents for the cations named above, while the F content was determined by wet chemical analysis (these specimens are denoted by ** under Analytical Method). The weight oxide percents of the cations for the remaining five specimens were determined by energy dispersive X-ray analysis (EDAX) with the F content being determined by wet chemical analysis (these specimens are denoted by + under Analytical Method). The number of OH ions per molecule was calculated on the assumption that the anion total (F + Cl + OH) in each molecule equalled 2.00. ND in the individual analyses indicates that the oxide percent for that element was not determined.

Table 2a.
Index of Specimens for Amphibole Analyses

| Analysis | Sample # [†] | Mineral Species | Analytical |
|----------|-----------------------|-----------------------|------------|
| | | | Method |
| 1 | A-77 | Tremolite | * |
| 2 | A-32 | Tremolite | + |
| 3 | A-149 | Tremolite | + |
| 4 | A-159 | Tremolite | * |
| 5 | A-116 | Tremolitic Hornblende | + |
| 6 | A-136 | Tremolitic Hornblende | * |
| 7 | A-60 | Edenite | * |
| 8 | A-84 | Edenite | + |
| 9 | A-15 | Edenite | * |
| 10 | A-29 | Edenite | ** |
| 11 | A-28 | Pargasitic Hornblende | ** |
| 12 | A-11 | Pargasitic Hornblende | ** |
| 13 | A-121 | Pargasite | * |
| 14 | A-26 | Pargasite | + |
| 15 | A-38 | Pargasite | ** |
| 16 | A-89 | Pargasite | * |

[†]Indicates mineral locality (See Figure 1)

Table 2b.
Analyses of Amphiboles

| | #1 | #2 | #3 | #4 | #5 | #6 |
|--------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|
| SiO ₂ | 56.00 | 56.23 | 55.72 | 56.47 | 53.90 | 53.90 |
| Al ₂ O ₃ | 1.18 | 1.51 | 1.47 | 2.26 | 3.43 | 4.20 |
| TiO ₂ | 0.06 | <0.05 | <0.05 | 0.06 | 0.10 | 0.17 |
| FeO | 1.00 | 0.45 | 1.34 | 0.43 | 0.30 | 0.81 |
| MgO | 23.38 | 23.72 | 23.48 | 23.89 | 23.72 | 23.23 |
| Mn | 0.05 | ND | ND | 0.03 | ND | 0.06 |
| CaO | 13.60 | 13.20 | 13.02 | 13.75 | 13.64 | 13.90 |
| Na ₂ O | 1.21 | 1.62 | 1.73 | 0.97 | 1.57 | 1.22 |
| K ₂ O | 0.16 | 0.05 | 0.05 | 0.41 | 0.15 | 0.38 |
| H ₂ O ⁺ | 1.28 | 1.43 | 1.84 | 1.47 | 0.93 | 1.45 |
| F | 1.80 | 1.55 | 0.69 | 1.52 | 2.61* | 1.54 |
| Cl | <u>0.09</u> | <u>ND</u> | <u>ND</u> | <u>0.03</u> | <u>ND</u> | <u>0.02</u> |
| Total | 99.81 | 99.76 | 99.34 | 101.29 | 100.35 | 100.88 |
| F+Cl=0 | <u>0.78</u> | <u>0.65</u> | <u>0.29</u> | <u>0.65</u> | <u>1.10</u> | <u>0.65</u> |
| Corrected Total | 99.03 | 99.11 | 99.05 | 100.64 | 99.25 | 100.23 |

Ions per formula unit based on 23 oxygen equivalents

| | | | | | | |
|-----------------|---------|------|------|------|------|------|
| Si | 7.77 | 7.76 | 7.73 | 7.69 | 7.48 | 7.42 |
| Al ⁴ | 0.19 | 0.24 | 0.24 | 0.31 | 0.52 | 0.58 |
| Al ⁶ | -- | 0.01 | -- | 0.05 | 0.04 | 0.10 |
| Ti | 0.01 | -- | -- | 0.01 | 0.01 | 0.02 |
| Fe | 0.12 | 0.05 | 0.15 | 0.05 | 0.03 | 0.09 |
| Mg | 4.84 | 4.88 | 4.85 | 4.85 | 4.90 | 4.76 |
| Mn | 0.01 | -- | -- | -- | -- | 0.01 |
| Ca | 2.02 | 1.94 | 1.94 | 2.01 | 2.03 | 2.05 |
| Na | 0.16 | 0.43 | 0.47 | 0.26 | 0.42 | 0.33 |
| K | 0.01 | 0.01 | 0.01 | 0.07 | 0.03 | 0.07 |
| OH | 1.19 | 1.32 | 1.70 | 1.34 | 0.86 | 1.33 |
| F | 0.79 | 0.68 | 0.30 | 0.65 | 1.14 | 0.67 |
| Cl | 0.02 | -- | -- | 0.01 | -- | -- |
| Ratio F | .40 | .34 | .15 | .33 | .57 | .34 |
| | F+Cl+OH | | | | | |

*Average of two F analyses by wet chemical procedure.

| | #7 | #8 | #9 | #10 | #11 | #12 |
|--------------------------------|--------|--------|--------|--------|--------|-------|
| SiO ₂ | 53.70 | 50.86 | 51.55 | 50.55 | 44.78 | 42.75 |
| Al ₂ O ₃ | 4.02 | 6.75 | 6.66 | 6.90 | 13.44 | 13.66 |
| TiO ₂ | 0.20 | <0.05 | 0.32 | 0.51 | 0.52 | 0.51 |
| FeO | 0.37 | 5.41 | 0.41 | 1.30 | 3.54 | 6.87 |
| MgO | 23.70 | 18.72 | 22.74 | 22.06 | 18.35 | 15.54 |
| MnO | 0.02 | ND | 0.02 | 0.07 | 0.06 | 0.14 |
| CaO | 13.68 | 12.56 | 13.79 | 13.30 | 13.14 | 12.77 |
| Na ₂ O | 1.71 | 1.38 | 1.87 | 2.35 | 2.76 | 2.06 |
| K ₂ O | 0.43 | 1.09 | 0.58 | 0.54 | 0.91 | 2.22 |
| H ₂ O ⁺ | 1.18 | 0.87 | 1.20 | 1.27 | 0.92 | 1.43 |
| F | 2.09 | 2.60* | 2.02 | 1.86 | 2.38* | 1.31 |
| Cl | 0.02 | ND | 0.04 | ND | ND | ND |
| Total | 101.12 | 100.24 | 101.20 | 100.71 | 100.80 | 99.26 |
| F+Cl+O | 0.88 | 1.09 | 0.86 | 0.78 | 1.00 | 0.55 |
| Corrected Total | 100.24 | 99.15 | 100.34 | 99.93 | 99.80 | 98.71 |

Ions per formula unit based on 23 oxygen equivalents

| | | | | | | |
|-----------------|---------|------|------|------|------|------|
| Si | 7.39 | 7.24 | 7.12 | 7.05 | 6.37 | 6.28 |
| Al ⁴ | 0.61 | 0.76 | 0.88 | 0.95 | 1.63 | 1.72 |
| Al ⁶ | 0.04 | 0.37 | 0.20 | 0.18 | 0.62 | 0.65 |
| Ti | 0.02 | -- | 0.03 | 0.05 | 0.06 | 0.06 |
| Fe | 0.04 | 0.64 | 0.05 | 0.15 | 0.42 | 0.34 |
| Mg | 4.86 | 3.97 | 4.68 | 4.58 | 3.89 | 3.40 |
| Mn | -- | -- | -- | 0.01 | 0.01 | 0.02 |
| Ca | 2.02 | 1.92 | 2.04 | 1.99 | 2.00 | 2.01 |
| Na | 0.46 | 0.38 | 0.50 | 0.63 | 0.76 | 0.59 |
| K | 0.08 | 0.20 | 0.10 | 0.10 | 0.17 | 0.42 |
| OH | 1.09 | 0.83 | 1.11 | 1.18 | 0.87 | 1.40 |
| F | 0.91 | 1.17 | 0.88 | 0.82 | 1.13 | 0.60 |
| Cl | -- | -- | 0.01 | -- | -- | -- |
| Ratio F | .45 | .59 | .44 | .41 | .57 | .33 |
| | F+Cl+OH | | | | | |

* Average of two F analyses by wet chemical procedure.

| | #13 | #14 | #15 | #16 |
|--------------------------------|-------------|-------------|-------------|-------------|
| SiO ₂ | 42.58 | 43.11 | 41.24 | 41.80 |
| Al ₂ O ₃ | 14.33 | 15.05 | 15.02 | 17.13 |
| TiO ₂ | 1.45 | 1.24 | 1.96 | 0.66 |
| FeO | 3.05 | 1.13 | 6.97 | 2.20 |
| MgO | 18.65 | 20.03 | 14.61 | 18.51 |
| MnO | 0.05 | ND | 0.20 | 0.04 |
| CaO | 13.10 | 12.79 | 12.46 | 13.40 |
| Na ₂ O | 2.78 | 3.28 | 2.56 | 3.05 |
| K ₂ O | 0.63 | 0.19 | 1.81 | 0.45 |
| H ₂ O ⁺ | 1.52 | 1.34 | 1.00 | 0.94 |
| F | 1.12 | 1.63 | 2.19 | 2.45 |
| Cl | <u>0.14</u> | <u>ND</u> | <u>ND</u> | <u>0.04</u> |
| Total | 99.40 | 99.79 | 100.02 | 100.67 |
| F+Cl=0 | <u>0.50</u> | <u>0.69</u> | <u>0.92</u> | <u>1.32</u> |
| Corrected Total | 98.90 | 99.10 | 99.10 | 99.35 |

| | | | | |
|-------------------|------|------|------|------|
| Si | 6.11 | 6.10 | 6.05 | 5.94 |
| Al ⁴ | 1.89 | 1.90 | 1.95 | 2.06 |
| Al ⁶ | 0.54 | 0.61 | 0.65 | 0.81 |
| Ti | 0.16 | 0.13 | 0.22 | 0.07 |
| Fe | 0.37 | 0.13 | 0.86 | 0.26 |
| Mg | 3.99 | 4.22 | 3.19 | 3.92 |
| Mn | 0.01 | -- | 0.02 | 0.01 |
| Ca | 2.01 | 1.94 | 1.96 | 2.04 |
| Na | 0.77 | 0.90 | 0.73 | 0.84 |
| K | 0.12 | 0.03 | 0.34 | 0.08 |
| OH | 1.46 | 1.27 | 0.98 | 0.89 |
| F | 0.51 | 0.73 | 1.02 | 1.10 |
| Cl | 0.03 | -- | -- | 0.01 |
| Ratio F / F+Cl+OH | .26 | .39 | .51 | .55 |

Phlogopite

The phlogopite specimens were all analyzed by electron micro-probe which yielded weight percent oxides for the cations (Si, Al, Ti, Fe, Mg, Mn, Ca, Na, K) and for F and Cl. The number of OH ions per molecule was calculated on the assumption that the anion total (F+Cl+OH) equalled 4.00.

Table 3a.

Index of Specimens for Phlogopite Analyses

| Analysis | Sample #* | Mineral Species |
|----------|-----------|-----------------|
| 1 | P-7 | Phlogopite |
| 2 | P-157 | Phlogopite |
| 3 | P-18A | Phlogopite |
| 4 | P-200-44 | Phlogopite |
| 5 | P-15 | Phlogopite |
| 6 | P-77 | Phlogopite |
| 7 | P-27 | Phlogopite |
| 8 | P-153 | Phlogopite |
| 9** | P-84 | Phlogopite |
| 10 | P-18 | Phlogopite |
| 11 | P-28 | Phlogopite |
| 12 | P-11 | Phlogopite |
| 13 | P-462-44 | Phlogopite |
| 14 | P-121 | Phlogopite |
| 15 | P-97 | Phlogopite |
| 16 | P-94 | Phlogopite |
| 17 | P-1 | Phlogopite |
| 18 | P-55A | Phlogopite |
| 19 | P-74 | Phlogopite |
| 20 | P-29 | Phlogopite |
| 21 | P-128 | Phlogopite |
| 22 | P-26 | Phlogopite |

*Indicates mineral locality (See Figure 1).

**Average of two analyses.

Table 3b.

Analyses of Phlogopite

| | #1 | #2 | #3 | #4 | #5 | #6 |
|--------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|
| SiO ₂ | 43.18 | 43.09 | 42.78 | 42.25 | 42.53 | 42.39 |
| Al ₂ O ₃ | 12.24 | 12.75 | 12.60 | 11.56 | 12.93 | 13.25 |
| TiO ₂ | 0.13 | 0.04 | 0.06 | 0.21 | 0.04 | 0.18 |
| FeO | 1.17 | 0.75 | 1.19 | 0.79 | 2.55 | 0.74 |
| MgO | 26.35 | 27.84 | 27.42 | 27.96 | 26.51 | 27.23 |
| MnO | 0.02 | 0.01 | 0.01 | 0.02 | 0.03 | 0.01 |
| CaO | 0.01 | 0.01 | 0.06 | 0.04 | 0.01 | <0.01 |
| Na ₂ O | 0.02 | 0.77 | 0.25 | 0.50 | 0.44 | 0.67 |
| K ₂ O | 10.23 | 9.26 | 10.00 | 8.90 | 9.89 | 9.20 |
| H ₂ O ⁺ | 2.41 | 2.77 | 2.21 | 2.48 | 1.95 | 2.23 |
| F | 3.79 | 3.21 | 4.31 | 3.60 | 4.82 | 4.23 |
| Cl | <u>0.03</u> | <u>0.02</u> | <u>0.03</u> | <u>0.07</u> | <u>0.11</u> | <u>0.06</u> |
| Total | 99.58 | 100.52 | 100.92 | 99.38 | 101.81 | 100.19 |
| F+Cl=0 | <u>1.60</u> | <u>1.36</u> | <u>1.82</u> | <u>1.53</u> | <u>2.05</u> | <u>1.79</u> |
| Corrected Total | 97.98 | 99.16 | 99.10 | 97.85 | 99.76 | 98.40 |

Ions per formula unit based on 22 oxygen equivalents

| | | | | | | |
|---------------------------|------|------|------|------|------|------|
| Si | 6.14 | 6.02 | 6.01 | 6.01 | 5.98 | 5.97 |
| Al ⁴ | 1.86 | 1.98 | 1.99 | 1.94 | 2.02 | 2.03 |
| Al ⁶ | 0.19 | 0.12 | 0.10 | -- | 0.12 | 0.17 |
| Ti | 0.01 | -- | 0.01 | 0.02 | -- | 0.02 |
| Fe | 0.14 | 0.09 | 0.14 | 0.21 | 0.30 | 0.09 |
| Mg | 5.58 | 5.78 | 5.74 | 5.93 | 5.56 | 5.72 |
| Mn | -- | -- | -- | -- | -- | -- |
| Ca | -- | -- | 0.01 | 0.01 | -- | -- |
| Na | -- | 0.21 | 0.07 | 0.14 | 0.12 | 0.18 |
| K | 1.85 | 1.65 | 1.79 | 1.62 | 1.77 | 1.65 |
| OH | 2.29 | 2.58 | 2.07 | 2.36 | 1.83 | 2.11 |
| F | 1.70 | 1.42 | 1.92 | 1.62 | 2.14 | 1.88 |
| Cl | 0.01 | -- | 0.01 | 0.02 | 0.03 | 0.01 |
| Ratio $\frac{F}{F+Cl+OH}$ | .43 | .36 | .48 | .41 | .54 | .47 |

| | #7 | #8 | #9 * | #10 | #11 * | #12 |
|--------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|
| SiO ₂ | 42.62 | 42.10 | 42.07 | 41.31 | 41.58 | 40.97 |
| Al ₂ O ₃ | 13.33 | 13.40 | 13.38 | 13.11 | 13.62 | 14.13 |
| TiO ₂ | 0.19 | 0.36 | 0.44 | 0.47 | 0.31 | 0.70 |
| FeO | 1.26 | 0.80 | 3.94 | 6.02 | 3.61 | 5.81 |
| MgO | 27.28 | 26.99 | 25.23 | 23.58 | 25.27 | 23.02 |
| MnO | 0.01 | 0.01 | 0.03 | 0.05 | 0.04 | 0.07 |
| CaO | 0.01 | 0.02 | <0.01 | <0.01 | 0.01 | 0.01 |
| Na ₂ O | 0.65 | 0.20 | 0.41 | 0.19 | 0.69 | 0.46 |
| K ₂ O | 9.51 | 10.11 | 9.93 | 10.03 | 9.48 | 9.49 |
| K ₂ O ⁺ | 2.34 | 2.26 | 2.39 | 2.46 | 2.49 | 2.74 |
| F | 4.09 | 4.16 | 3.58 | 3.48 | 3.60 | 2.99 |
| Cl | <u>0.04</u> | <u>0.09</u> | <u>0.05</u> | <u>0.27</u> | <u>0.11</u> | <u>0.12</u> |
| Total | 101.33 | 100.50 | 101.60 | 100.97 | 100.81 | 100.51 |
| F+Cl=0 | <u>1.73</u> | <u>1.77</u> | <u>1.52</u> | <u>1.53</u> | <u>1.54</u> | <u>1.29</u> |
| Corrected Total | 99.60 | 98.73 | 100.08 | 99.44 | 99.27 | 99.22 |

Ions per formula unit based on 22 oxygen equivalents

| | | | | | | |
|-----------------|------|------|------|------|------|------|
| Si | 5.95 | 5.94 | 5.93 | 5.93 | 5.90 | 5.86 |
| Al ⁴ | 2.05 | 2.06 | 2.07 | 2.07 | 2.10 | 2.14 |
| Al ⁶ | 0.14 | 0.16 | 0.15 | 0.15 | 0.18 | 0.24 |
| Ti | 0.02 | 0.04 | 0.05 | 0.05 | 0.03 | 0.07 |
| Fe | 0.15 | 0.09 | 0.46 | 0.72 | 0.43 | 0.70 |
| Mg | 5.68 | 5.67 | 5.30 | 5.04 | 5.34 | 4.91 |
| Mn | -- | -- | -- | 0.01 | 0.01 | 0.01 |
| Ca | -- | -- | -- | -- | -- | -- |
| Na | 0.18 | 0.05 | 0.11 | 0.05 | 0.20 | 0.13 |
| K | 1.70 | 1.82 | 1.78 | 1.84 | 1.71 | 1.73 |
| OH | 2.18 | 2.13 | 2.39 | 2.36 | 2.36 | 2.62 |
| F | 1.81 | 1.85 | 1.60 | 1.58 | 1.61 | 1.35 |
| Cl | 0.01 | 0.02 | 0.01 | 0.06 | 0.03 | 0.03 |
| Ratio F | .45 | .46 | .40 | .40 | .40 | .34 |

* F+Cl+OH

* Average of two analyses.

| | #13 | #14 | #15 | #16 | #17 | #18 |
|--------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|
| SiO ₂ | 41.14 | 41.65 | 39.78 | 41.46 | 41.20 | 41.12 |
| Al ₂ O ₃ | 14.19 | 14.25 | 13.92 | 14.44 | 15.16 | 15.32 |
| TiO ₂ | 0.52 | 0.61 | 0.88 | 0.92 | 0.14 | 0.17 |
| FeO | 4.06 | 2.30 | 8.03 | 1.85 | 1.00 | 0.60 |
| MgO | 24.48 | 26.15 | 21.26 | 26.30 | 26.97 | 27.44 |
| MnO | 0.06 | <0.01 | 0.12 | 0.02 | 0.01 | <0.01 |
| CaO | 0.02 | <0.01 | 0.01 | <0.01 | 0.01 | 0.02 |
| Na ₂ O | 0.40 | 1.04 | 0.28 | 1.04 | 1.19 | 1.72 |
| K ₂ O | 9.83 | 9.00 | 9.63 | 8.97 | 8.98 | 7.71 |
| H ₂ O ⁺ | 2.49 | 3.13 | 2.22 | 3.35 | 2.80 | 2.62 |
| F | 3.56 | 2.37 | 3.88 | 1.94 | 2.82 | 3.51 |
| Cl | <u>0.12</u> | <u>0.09</u> | <u>0.18</u> | <u>0.05</u> | <u>0.06</u> | <u>0.06</u> |
| Total | 100.87 | 100.59 | 100.19 | 100.34 | 100.33 | 100.29 |
| F+Cl=0 | <u>1.53</u> | <u>1.02</u> | <u>1.67</u> | <u>0.83</u> | <u>1.20</u> | <u>1.49</u> |
| Corrected Total | 99.34 | 99.57 | 98.52 | 99.51 | 99.13 | 98.80 |


Ions per formula unit based on 22 oxygen equivalents

| | | | | | | |
|--------------------|------|------|------|------|------|------|
| Si | 5.85 | 5.83 | 5.82 | 5.80 | 5.76 | 5.74 |
| Al ⁴ | 2.15 | 2.17 | 2.18 | 2.20 | 2.24 | 2.26 |
| Al ⁶ | 0.23 | 0.18 | 0.22 | 0.18 | 0.21 | 0.26 |
| Ti | 0.05 | 0.06 | 0.10 | 0.10 | 0.01 | 0.02 |
| Fe | 0.48 | 0.27 | 0.98 | 0.22 | 0.12 | 0.07 |
| Mg | 5.19 | 5.46 | 4.63 | 5.48 | 5.62 | 5.71 |
| Mn | 0.01 | -- | 0.01 | -- | -- | -- |
| Ca | -- | -- | -- | -- | -- | -- |
| Na | 0.11 | 0.28 | 0.08 | 0.28 | 0.32 | 0.46 |
| K | 1.78 | 1.61 | 1.80 | 1.60 | 1.60 | 1.37 |
| OH | 2.37 | 2.93 | 2.17 | 3.13 | 2.61 | 2.44 |
| F | 1.60 | 1.05 | 1.79 | 0.86 | 1.25 | 1.55 |
| Cl | 0.03 | 0.02 | 0.04 | 0.01 | 0.14 | 0.01 |
| Ratio F F+Cl+OH | .40 | .26 | .45 | .22 | .32 | .39 |

| | #19 | #20 | #21 | #22 |
|--------------------------------|-------------|-------------|-------------|-------------|
| SiO ₂ | 41.15 | 40.29 | 39.96 | 38.96 |
| Al ₂ O ₃ | 15.19 | 16.10 | 16.33 | 17.46 |
| TiO ₂ | 0.03 | 1.00 | 1.21 | 1.64 |
| FeO | 0.23 | 1.15 | 2.87 | 0.93 |
| MgO | 28.33 | 26.17 | 25.09 | 25.49 |
| MnO | 0.01 | 0.02 | 0.03 | 0.02 |
| CaO | 0.08 | 0.02 | 0.01 | 0.02 |
| Na ₂ O | 1.92 | 1.09 | 1.66 | 1.34 |
| K ₂ O | 7.07 | 8.93 | 7.80 | 8.53 |
| H ₂ O ⁺ | 2.67 | 2.92 | 3.20 | 3.39 |
| F | 3.43 | 2.81 | 2.21 | 1.79 |
| Cl | <u>0.05</u> | <u>0.13</u> | <u>0.12</u> | <u>0.13</u> |
| Total | 100.16 | 100.63 | 100.49 | 99.70 |
| F+Cl=0 | <u>1.45</u> | <u>1.21</u> | <u>0.96</u> | <u>0.78</u> |
| Corrected Total | 98.71 | 99.42 | 99.53 | 98.92 |

Ions per formula unit based on 22 oxygen equivalents

| | | | | |
|--------------------|------|------|------|------|
| Si | 5.73 | 5.63 | 5.60 | 5.47 |
| Al ⁴ | 2.27 | 2.37 | 2.40 | 2.53 |
| Al ⁶ | 0.22 | 0.28 | 0.30 | 0.36 |
| Ti | -- | 0.10 | 0.13 | 0.17 |
| Fe | 0.03 | 0.13 | 0.34 | 0.11 |
| Mg | 5.88 | 5.46 | 5.24 | 5.33 |
| Mn | -- | -- | -- | -- |
| Ca | 0.01 | -- | -- | -- |
| Na | 0.52 | 0.29 | 0.45 | 0.36 |
| K | 1.26 | 1.59 | 1.39 | 1.53 |
| OH | 2.48 | 2.73 | 2.99 | 3.18 |
| F | 1.51 | 1.24 | 0.98 | 0.79 |
| Cl | 0.01 | 0.03 | 0.03 | 0.03 |
| Ratio F F+Cl+OH | .38 | .31 | .25 | .20 |



LOCATION MAP - FIGURE 1