

## **Kiglapait mineralogy V: Feldspars in a hot, dry magma**

**S.A. MORSE<sup>1,\*</sup>**

<sup>1</sup>Department of Geosciences, University of Massachusetts, 611 North Pleasant Street, Amherst, Massachusetts, 01003-9297, U.S.A.

### **ABSTRACT**

The lithology of the 1.305 Ga Kiglapait intrusion is dominated by a Lower Zone of troctolite, succeeded by an Upper Zone of olivine gabbro, ferrodiorite, and syenite with olivine composition of pure fayalite. The feldspar composition of the intrusion varies from An<sub>68</sub> to An<sub>9</sub> over a thickness of 8.4 km from the base to a sandwich horizon under an Upper Border Zone. The anhydrous nature of the Kiglapait syenites is shown by their high temperature, by the loss of minor biotite up-stratigraphy in the intrusion, and the absence of amphibole. The end-stage feldspar of the Kiglapait syenites is that of a solidus embedded in a solvus in a 3 kbar eutectic at 1000 °C. The end-member assemblage at temperature and pressure is invariant. The final bulk composition is relatively An-rich—An ~11%—with a composition of  $X_{Or} = 1/3$  when projected to the Ab-Or sideline. The experimental feldspar solvus when corrected for the effects of An and Ba and referred to 3 kbar penetrates the solidus and fits the experimental tie lines. These conditions precede a stage of local coarsening under subsolidus conditions that is found in colloform symplectites invading mesoperthite. The oligoclase-orthoclase symplectites are iso-compositional with their host mesoperthites. The coarsening is assumed to be related to a plausibly F-rich vapor phase that is locally consumed with time. The observed phase compositions indicate the end of exsolution at ~800 °C at 3 kbar on the binodal solvus.

**Keywords:** Feldspar compositions, Kiglapait Intrusion, chemistry, textures, exsolution, symplectite, coarsening, solvus, syenites, cooling history