

***P-V-T* equation of state of hydrous phase A up to 10.5 GPa**

CUIPING YANG¹, TORU INOUE^{1,2,*},†, AND TAKUMI KIKEGAWA³

¹Geodynamics Research Center, Ehime University, Matsuyama, Ehime 790-8577, Japan

²Department of Earth and Planetary Systems Science & Hiroshima Institute of Plate Convergence Region Research (HiPeR), Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8526, Japan

³Photon Factory, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan

ABSTRACT

Pressure-volume-temperature (*P-V-T*) data of synthetic Mg₇Si₂O₈(OH)₆ phase A were collected under *P-T* conditions up to ~10.5 GPa and 900 K by energy-dispersive X-ray diffraction using a cubic type multi-anvil apparatus, MAX80, located at the Photon Factory–Advanced Ring (PF-AR) at the High Energy Accelerator Research Organization (KEK). *P-V* EoS using only room-temperature data yielded $V_0 = 511.6(2) \text{ \AA}^3$, $K_{T0} = 106.8(18) \text{ GPa}$, and pressure derivative $K'_T = 3.88(38)$. These parameters were consistent with the subsequent equation of state (EoS) analysis. The compressibility of phase A was anisotropic, with its *a*-axis being ~26% more compressible than the *c*-axis, which is normal to the plane of the distorted close-packed layers. A fit of the present data to the high-temperature Birch-Murnaghan EoS yielded $V_0 = 511.7(3) \text{ \AA}^3$, $K_0 = 104.4(24) \text{ GPa}$, $K' = 4.39(48)$, $(\partial K_T / \partial T)_P = -0.027(5) \text{ GPa K}^{-1}$, and thermal expansion $\alpha = a + bT$ with values of $a = 2.88(27) \times 10^{-5} \text{ K}^{-1}$ and $b = 3.54(68) \times 10^{-8} \text{ K}^{-2}$. The lattice dynamical approach by the Mie-Grüneisen-Debye EoS yielded $\theta_0 = 928(114) \text{ K}$, $q = 2.9(10)$, and $\gamma_0 = 1.19(8)$. The isobaric heat capacity C_p of phase A at 1 atm. was calculated based on the Mie-Grüneisen-Debye EoS fit of present *P-V-T* data. In addition, the density profiles of subducting slabs with different degrees of serpentinization were also calculated along the cold geotherm up to ~13 GPa. The serpentinization of subducting slab will significantly lower the density of slab at shallower depth; however, this effect becomes negligible when antigorite dehydrates to phase A. Because the phase A bearing subducting slab is supposed to be denser than the surrounding mantle, the water can transport into deeper parts of the upper mantle and the transition zone.

Keywords: Bulk modulus, phase A, dense hydrous magnesium silicate (DHMS), equation of state (EoS), heat capacity, high pressure and high temperature (HPHT), thermal expansion; Physics and Chemistry of Earth's Deep Mantle and Core