

## Discovery of terrestrial andreyivanovite, FeCrP, and the effect of Cr and V substitution on the low-pressure barringerite-allabogdanite transition

EVGENY V. GALUSKIN<sup>1,\*</sup>, JOACHIM KUSZ<sup>2</sup>, IRINA O. GALUSKINA<sup>1,†</sup>, MARIA KSIĄŻEK<sup>2</sup>,  
YEVEGENY VAPNIK<sup>3</sup>, AND GRZEGORZ ZIELIŃSKI<sup>4</sup>

<sup>1</sup>Faculty of Natural Sciences, University of Silesia, Będzińska 60, 41-200 Sosnowiec, Poland

<sup>2</sup>Faculty of Science and Technology, University of Silesia, ul. 75. Pułku Piechoty 1, 41-500 Chorzów, Poland

<sup>3</sup>Department of Geological and Environmental Sciences, Ben-Gurion University of the Negev, P.O.B. 653, Beer-Sheva 84105, Israel

<sup>4</sup>Micro-Area Analysis Laboratory, Polish Geological Institute—National Research Institute, Rakowiecka 4, 00-975 Warsaw, Poland

### ABSTRACT

Iron phosphides with significant variations of Cr (up to 18 wt%) and V (up to 8.6 wt%) contents were detected in gehlenite-bearing breccia at the Hatrurim Complex, Negev desert, Israel. Investigations of the composition and structure of the Fe<sub>2</sub>P phosphides showed that when the V+Cr content is higher than 0.26 apfu (atoms per formula unit), a transition from the hexagonal barringerite (*P6̄2m*) to orthorhombic allabogdanite (*Pnma*) takes place. According to the experimental data, allabogdanite is a high-pressure (>8 GPa) polymorph of barringerite. Pseudowollastonite associated with Cr-V-bearing allabogdanite is an indicator of phosphide crystallization at high temperature (>1200 °C) and low pressure. Thus, at the low pressure close to ambient, when more than 13 at% Fe in Fe<sub>2</sub>P is substituted by Cr and V, the orthorhombic polymorph is stable. The orthorhombic phosphide with the highest Cr and V contents belongs to the andreyivanovite species with the FeCrP end-member formula. This is the first finding on Earth of that very rare mineral described from the Kaidun meteorite. Some Cr-V-bearing phosphides have an unusual morphology, which cannot be explained by crystallization from a melt. More probably, these phosphides can form in the process of replacing fish bone remains. We believe that sedimentary protolith was not thermally altered and contained a significant amount of bituminous organic matter and phosphorite inclusions. Injecting paralava into the sedimentary rocks determines the conditions for phosphide formation on the boundary of these rocks as a result of the high-temperature carbothermal reduction process.

**Keywords:** Terrestrial natural phosphides, barringerite, allabogdanite, andreyivanovite, phase transition, Hatrurim Complex