

## **Synthesis of boehmite-type GaOOH: A new polymorph of Ga oxyhydroxide and geochemical implications**

**MENG LIU<sup>1,2</sup>, WEI YIN<sup>1,2</sup>, HAO-FAN JIANG<sup>1,2</sup>, YU-HAN WANG<sup>1,2</sup>, QI-TAO HU<sup>1,2</sup>, TIAN-LEI ZHAO<sup>1,2</sup>,  
QI-ZHI YAO<sup>3,\*</sup>, SHENG-QUAN FU<sup>4</sup>, AND GEN-TAO ZHOU<sup>1,2,5,\*</sup>**

<sup>1</sup>Deep Space Exploration Laboratory/School of Earth and Space Sciences, University of Science and Technology of China, Hefei 230026, China

<sup>2</sup>CAS Key Laboratory of Crust-Mantle Materials and Environments, School of Earth and Space Sciences, University of Science and Technology of China, Hefei 230026, China

<sup>3</sup>School of Chemistry and Materials Science, University of Science and Technology of China, Hefei 230026, China

<sup>4</sup>Hefei National Laboratory for Physical Sciences at the Microscale, University of Science and Technology of China, Hefei 230026, China

<sup>5</sup>CAS Center for Excellence in Comparative Planetology, Hefei 230026, China

### **ABSTRACT**

Gallium (Ga) and aluminum (Al) belong to group IIIA elements in the periodic table. They show a coupled geochemical behavior in most natural systems and are considered as “geochemical partners.” However, compared with the principal oxyhydroxides of Al in nature, gibbsite [Al(OH)<sub>3</sub>], boehmite ( $\gamma$ -AlOOH), and diaspore ( $\alpha$ -AlOOH), only the analogs söhngeite [Ga(OH)<sub>3</sub>] and tsumgallite ( $\alpha$ -GaOOH) were reported. In this work, boehmite-type GaOOH ( $\gamma$ -GaOOH), a new polymorph of GaOOH, was synthesized for the first time using boehmite ( $\gamma$ -AlOOH) as a template. The synthesized  $\gamma$ -GaOOH was characterized by a series of techniques, including X-ray diffraction (XRD), high-angle annular dark-field scanning transmission electron microscopy (HAADF-STEM), and selected area electron diffraction (SAED). Furthermore, a model based on the boehmite structure was successfully applied to define the  $\gamma$ -GaOOH structure by the Rietveld method. Results from sample characterization and structural refinement support the successful synthesis of boehmite-type GaOOH, and thus it is referred to as  $\gamma$ -GaOOH. The synthesis of  $\gamma$ -GaOOH in the laboratory is valuable to understanding the Ga geochemistry and its enrichment process in Ga-rich boehmite in coal and bauxite.

**Keywords:** Gallium, boehmite analog,  $\gamma$ -GaOOH, template synthesis, Ga geochemistry