

Crystal structure of kanemite, $\text{NaHSi}_2\text{O}_5 \cdot 3\text{H}_2\text{O}$, from the Aris phonolite, Namibia

**LAURENCE A.J. GARVIE,^{1,*} BERTRAND DEVOUARD,^{1,†} THOMAS L. GROY,² FERNANDO CÁMARA,^{1,‡}
AND PETER R. BUSECK^{1,2}**

¹Department of Geology, Arizona State University, Tempe, Arizona 85287-1404, U.S.A.

²Department of Chemistry/Biochemistry, Arizona State University, Tempe, Arizona 85287, U.S.A.

ABSTRACT

Kanemite was studied by single-crystal X-ray diffraction. The mineral, ideally $\text{NaHSi}_2\text{O}_5 \cdot 3\text{H}_2\text{O}$, is orthorhombic (space group *Pbcn*); unit-cell parameters are $a = 4.946(3)$, $b = 20.502(15)$, $c = 7.275(3)$ Å, with $Z = 4$. The structure is solved and refined to an R value of 0.058 for 825 independent reflections. The arrangement of atoms consists of alternating (010) sheets of corrugated $[\text{Si}_2\text{O}_4\text{OH}]_n^-$ and hydrated Na. The silicate sheets contain six-membered rings of $\text{HOSiO}_3\text{-SiO}_4$ units. Sodium atoms coordinate to six water molecules, forming layers of distorted octahedra. Residual electron densities were located that give reasonable positions for four H atoms. One H is part of a silanol group, and the other three H atoms are associated with water bonded to Na. Bonding between the silicate and Na sheets is through hydrogen bonding from H of the Na layer to O of the silicate sheet.