

## **Two types of metamorphic monazite with contrasting La/Nd, Th, and Y signatures in an ultrahigh-pressure metapelite from the Pohorje Mountains, Slovenia: Indications for pressure-dependent REE exchange between apatite and monazite?**

**ERWIN KRENN,<sup>1,\*</sup> MARIAN JANÁK,<sup>2</sup> FRITZ FINGER,<sup>1</sup> IGOR BROSKA,<sup>2</sup> AND PATRIK KONEČNÝ<sup>3</sup>**

<sup>1</sup>Department of Materials Engineering and Physics, University of Salzburg, Hellbrunnerstrasse 34, 5020 Salzburg, Austria

<sup>2</sup>Geological Institute, Slovak Academy of Sciences, Dúbravská cesta 9, 840 05 Bratislava, Slovak Republic

<sup>3</sup>Dionýz Štúr Institute of Geology, Mlynská dolina 1, 817 04 Bratislava, Slovak Republic

### **ABSTRACT**

Two monazite generations (M1; M2) were distinguished in a kyanite-garnet gneiss from the UHP terrain of the Pohorje Mountains, Slovenia. *P-T* estimates reveal a peak event at 760 °C/2.6 GPa and isothermal decompression down to 700 °C/0.6 GPa.

M1 type provides a Th-U-Pb mean date of  $100 \pm 6$  Ma, ThO<sub>2</sub> contents between 3–7 wt%, Y<sub>2</sub>O<sub>3</sub> values <0.3 wt%, and La/Nd ratios (1.2–1.4) that are clearly higher than for the whole-rock La/Nd (1.1). The absence of Y zoning in M1 and the lack of monazite inclusions in garnet indicate that M1 formed after the main stage of garnet growth (>1.2 MPa), probably close to the *P-T* peak.

M2 type is slightly younger than M1 ( $74 \pm 16$  Ma), and has a lower La/Nd (0.3–0.9), lower ThO<sub>2</sub> (0.1–5 wt%), and higher Y<sub>2</sub>O<sub>3</sub> (up to 3.2 wt%). Most M2 monazites occur as tiny needles within apatite (subtype M2-a) or along apatite margins (M2-b). Parasitic growth of M2-a and -b from apatite is supported by its low ThO<sub>2</sub> (<1 wt%) and La/Nd (<0.5). Isolated matrix grains (M2-c) and overgrowths around M1 (M2-d) have slightly higher La/Nd (0.5–0.9) and higher ThO<sub>2</sub> (5 wt%) and were supplied from an apatite and M1 source. Elevated yttrium suggests that M2 formed during decompression, when garnet was consumed and Y was released.

These observations imply that at UHP conditions MREE-rich apatite coexisted with low-MREE M1 monazite and reacted during decompression to Ca-F-apatite plus MREE-rich M2 monazite. This provides strong arguments that REE-partitioning between apatite and monazite is pressure-dependent.

**Keywords:** Monazite, apatite, REE distribution, ultra-high pressure, Pohorje Mountains