

LETTER

**Si-Al distribution in high-pressure  $\text{CaAl}_4\text{Si}_2\text{O}_{11}$  phase: A  $^{29}\text{Si}$  and  $^{27}\text{Al}$  NMR study**

**XIANYU XUE,<sup>1,\*</sup> SHUANGMENG ZHAI,<sup>2</sup> AND MASAMI KANZAKI<sup>1</sup>**

<sup>1</sup>Institute for Study of the Earth's Interior, Okayama University, Misasa, Tottori, 682-0193 Japan

<sup>2</sup>School of Earth and Space Sciences, Peking University, Beijing, 100871, P.R. China

**ABSTRACT**

High-resolution  $^{29}\text{Si}$  and  $^{27}\text{Al}$  NMR techniques have been applied to resolve the Si-Al distribution and coordination in the high-pressure  $\text{CaAl}_4\text{Si}_2\text{O}_{11}$  (CAS) phase, a potentially important mineral in subducted crustal materials in the deep mantle that has a unique hexagonal ferrite structure containing two octahedral (M1; M2) and one trigonal bipyramidal sites. The  $^{29}\text{Si}$  MAS NMR spectra of the CAS phase synthesized at 20 GPa and 1400~1600 °C show two broad, asymmetric peaks near -92.7 and -182.7 ppm with an intensity ratio of 1:3, suggesting that 1/4 of the Si are in tetrahedral coordination and 3/4 in octahedral coordination. Therefore, the trigonal bipyramidal and M1 octahedral sites are each occupied by equal proportions of Si and Al, and the former are effectively half-occupied face-sharing tetrahedra (at least for Si). The  $^{27}\text{Al}$  MAS and 3Q MAS NMR spectra contain only one unresolved peak typical of octahedral Al with a range of quadrupolar coupling constants.

**Keywords:** NMR spectroscopy, Al and Si coordination, CAS phase, Si-Al disorder, high pressure