

Sound velocities across calcite phase transitions by Brillouin scattering spectroscopy

CHAO-SHUAI ZHAO^{1,2}, HE-PING LI^{1,*}, PO-FEI CHEN³, AND JIAN-JUN JIANG¹

¹Key Laboratory of High-Temperature and High-Pressure Study of the Earth's Interior, Institute of Geochemistry, Chinese Academy of Sciences, 550081 Guiyang, China

²College of Earth Sciences, University of Chinese Academy of Sciences, 100049 Beijing, China

³Department of Earth Sciences, National Central University, 32001 Taoyuan, Taiwan, R.O.C.

ABSTRACT

Calcite (CaCO_3) is widely considered an important carbon carrier in the Earth's interior. Laboratory measurements of the velocities and elastic properties of calcite are important for understanding the deep carbon cycle. The sound velocities of calcite were determined up to 10.3 GPa at ambient temperature by Brillouin scattering spectroscopy. Dramatic decreases in the velocity of compressional wave (V_p) and shear wave (V_s) and abrupt increases in the V_p anisotropy (A_p) and maximum V_s anisotropy ($A_{s_{\max}}$) were detected across the phase transition from CaCO_3 -I to CaCO_3 -II. Dramatic increases in the V_p and V_s and an abrupt decrease in A_p were observed across the phase transition from CaCO_3 -II to CaCO_3 -III. The phase transition from CaCO_3 -I to CaCO_3 -II may potentially explain the Gutenberg discontinuity at 51 km in the Izu-Bonin region. The V_p and V_s values of calcite were low. Our new results combined with literature data suggest that the low velocities of CaCO_3 could potentially explain the low-velocity zone occurring in northeastern (NE) Japan.

Keywords: Brillouin scattering, sound velocity, elasticity, CaCO_3 , high pressure