

**High-pressure and high-temperature vibrational properties and anharmonicity of
carbonate minerals up to 6 GPa and 500 °C by Raman spectroscopy**

Stefan Farsang^{1*}, Remo N. Widmer² and Simon A. T. Redfern³

¹Department of Earth Sciences, University of Cambridge, Downing Street, Cambridge, CB2
3EQ, UK (*sf571@cam.ac.uk)

²Empa, Swiss Federal Laboratories for Materials Science and Technology, Laboratory for
Mechanics of Materials and Nanostructures, Feuerwerkerstrasse 39, Thun, 3602, Switzerland

³Asian School of the Environment, Nanyang Technological University, 50 Nanyang Avenue,
Singapore, 639798, Singapore

Supplementary material

Figure S1. Evolution of frequencies and full width half maxima (FWHM) of bands of the ν_1 internal vibrational mode in aragonite-group carbonate minerals as a function of pressure.




Figure S2. Evolution of frequencies and full width half maxima (FWHM) of bands of T and L external vibrational modes and ν_1 internal vibrational mode in calcite-group carbonate minerals as a function of pressure.




Figure S3. Evolution of frequencies and full width half maxima (FWHM) of bands of the ν_1 internal vibrational mode in aragonite-group carbonate minerals as a function of temperature.




Figure S4. Phase transition of aragonite to calcite recorded by Raman spectra taken at 425 °C and 450 °C.




Figure S5. Thermal decomposition of cerussite recorded by Raman spectra taken at 225 °C and 250 °C.




Figure S6. Evolution of frequencies and full width half maxima (FWHM) of bands of T and L external vibrational modes and ν_1 internal vibrational mode in calcite-group carbonate minerals as a function of temperature.




Figure S7. Thermal decomposition of gaspeite recorded by Raman spectra taken at 375 °C and 400 °C.




Figure S8. Thermal decomposition of smithsonite recorded by Raman spectra taken at 325 °C, 350 °C, 375 °C, and 400 °C.




Figure S9. a) isobaric Grüneisen parameters (γ_{iP}), b) isothermal Grüneisen parameters (γ_{iT}), and c) intrinsic anharmonic parameters (a_i) of the Raman-active vibrational modes of aragonite-group carbonate minerals as a function of band frequencies.




Figure S10. a) isobaric Grüneisen parameters (γ_{iP}), b) isothermal Grüneisen parameters (γ_{iT}), and c) intrinsic anharmonic parameters

(a) of the Raman-active vibrational modes of calcite-group carbonate minerals as a function of band frequencies.

