

Supplementary Material

Characterization and assessment of the potential toxicity/pathogenicity of Russian commercial chrysotile

DARIO DI GIUSEPPE^{1,2,*}, ALESSANDRO ZOBOLI¹, LUCA NODARI³, LUCA PASQUALI⁴, ORIETTA SALA¹, PAOLO BALLIRANO⁵, DANIELE Malferrari¹, SIMONA RANERI⁶, MIRIAM HANUSKOVA⁴, AND ALESSANDRO F. GUALTIERI¹

¹Department of Chemical and Geological Sciences, University of Modena and Reggio Emilia, Via G. Campi 103, Modena, I-41125, Italy.

²Department of Sciences and Methods for Engineering, University of Modena and Reggio Emilia, Via Amendola 2, Reggio Emilia, I-42122, Italy.

³ICMATE-CNR, Institute of Condensed Matter Chemistry and Technologies for Energy, Italian National Research Council, Corso Stati Uniti, 4, I-35127, Padova, Italy.

⁴Department of Engineering "Enzo Ferrari", University of Modena and Reggio Emilia, Via Pietro Vivarelli, 10, I-41125, Modena, Italy.

⁵Department of Earth Sciences, Sapienza University of Rome, Piazzale Aldo Moro 5, I-00185, Rome, Italy.

⁶ICCOM-CNR, Institute of Chemistry of Organometallic Compounds, Italian National Research Council, Via G. Moruzzi, 1, I-56124, Pisa, Italy.

* Corresponding author, e-mail address: ddigiuse@unimore.it, Tel. +390592058497




Figure OM1. Backscattered-Electron (BSE) images of chrysotile particles embedded in epoxy resin, acquired during the EPMA session. Large chrysotile bundles and clusters (**a**) and (**b**). Some fibers curl after being ripped from the bundle due to the sample lapping (**a**). Magnetite particles appear as bright white spots (**a**) and (**b**).




Figure OM2. EDX spectra acquired during the SEM investigation. **(a)** EDX spectrum of a chrysotile fiber bundle. **(b)** EDX spectrum of a magnetite particle.




Figure OM3. ^{57}Fe Mössbauer spectrum of Russian chrysotile fitted by the superimposition of three doublets (see text for details).




Figure OM4. FTIR spectrum of the Russian chrysotile (500–4000 cm^{-1}). For comparison, the FTIR trace of UICC standard chrysotile is included.




Figure OM5. The XPS Fe 2p region of the Russian chrysotile.

Table OM1. Mean values of the EPMA analyses of the Russian chrysotile. Wt% (weight percent); the standard deviation is shown in brackets. The chemical composition of the UICC chrysotile standard and other Italian chrysotile samples is reported for comparison (after Pollastri et al. 2016).

Wt%	Russian chrysotile	UICC chrysotile	Balangero chrysotile	Valmalenco chrysotile
SiO ₂	42.43 (1.5)	42.5(3)	40.6(5)	42.5(2)
TiO ₂	0.03 (0.01)	0.01(2)	0.01(2)	0.06(5)
Al ₂ O ₃	0.62 (0.2)	0.20(1)	2.40(6)	0.20(8)
Cr ₂ O ₃	0.14 (0.1)	0.05(4)	0.20(3)	0.08(8)
MnO	0.06 (0.01)	0.05(4)	0.06(4)	0.06(4)
MgO	41.21 (1.8)	41.9(2)	39.8(6)	41.6(2)
CaO	0.05 (0.01)	0.01(1)	0.02(2)	0.09(6)
Na ₂ O	0.02 (0.01)	0.013(9)	0.012(9)	0.02(1)
K ₂ O	0.01 (0.01)	0.004(5)	0.003(7)	0.05(9)
NiO	0.15 (0.1)	0.06(4)	0.05(7)	0.20(1)
FeO	0.70 (0.8)	1.40(3)	2.50(5)	1.40(1)
Fe ₂ O ₃	1.27 (0.8)	0.20(2)	0.40(3)	0.01(2)
Tot	86.67 (1.7)	86.4	86.1	86.3

References

- Pollastri, S., Perchiazzi, N., Lezzerini, M., Plaisier, J.R., Cavallo, A., Dalconi, M.C., Bursi Gandolfi, N., Gualtieri, A.F., 2016. The crystal structure of mineral fibres. 1. Chrysotile. *Periodico di Mineralogia*, 85, 249-259.