Identifying xenocrystic tourmaline in Himalayan leucogranites

JINSHENG HAN¹, HUAYONG CHEN^{2,*}, HAIJUN XU³, OLIVIER NADEAU⁴, AND CHANG XU³

¹State Key Laboratory of Geological Processes and Mineral Resources, and School of Earth Resources, China University of Geosciences, Wuhan (CUGW) 430074, China

²Key Laboratory of Mineralogy and Metallogeny, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China
³State Key Laboratory of Geological Processes and Mineral Resources, and School of Earth Sciences, CUGW, 430074, China
⁴Laboratory for Isotopes in Economic Geology and its Environment (LIEGE), Department of Mineral Resources and Engineering, School of Earth Sciences, CUGW, 430074, China

ABSTRACT

Tournaline is a common autocrystic mineral in granitic and pegmatitic rocks that records valuable information on the physicochemical evolution of magmas. However, xenocrystic tournaline that does not crystallize from host magmas is also common but seldom reported and notoriously difficult to identify. Here, autocrystic (Tur G) and xenocrystic (Tur P) tournalines are identified in two-mica granites and granitic pegmatites from the Cuonadong leucogranites, eastern Himalaya. Electron backscatter diffraction (EBSD), coupled with electron-probe micro-analysis (EPMA) and laser ablation ICP-MS analyses (LA-ICP-MS), is used to discriminate xenocrystic from autocrystic tournaline. Although both tournalines have slightly different chemical compositions, the differences are insufficient to permit unambiguous discrimination. However, EBSD systematically shows complex intra-crystalline orientations and deformation xenocrystic grains, and undeformed crystal lattices are predominant in autocrystic grains. EBSD could be a useful tool to identify the origin of tournaline in granitic rocks, particularly when geochemical analyses and/or geochronological measurements are inconclusive.

Keywords: Autocrystic and xenocrystic tourmaline, boron isotopes, Cuonadong leucogranites, electron backscattered diffraction analyses, Himalayan granites