

## Supporting Information

### **Synthesis of boehmite-type GaOOH: a new polymorph of Ga oxyhydroxide and geochemical implications**

Meng Liu<sup>a</sup>, Wei Yin<sup>a</sup>, Hao-Fan Jiang<sup>a</sup>, Yu-Han Wang<sup>a</sup>, Qi-Tao Hu<sup>a</sup>, Tian-Lei Zhao<sup>a</sup>, Qi-Zhi Yao<sup>b\*</sup>, Sheng-Quan Fu<sup>c</sup> and Gen-Tao Zhou<sup>a,d\*</sup>

<sup>a</sup> CAS Key Laboratory of Crust-Mantle Materials and Environments, School of Earth and Space Sciences, University of Science and Technology of China, Hefei 230026, P. R. China.

<sup>b</sup> School of Chemistry and Materials Science, University of Science and Technology of China, Hefei 230026, P. R. China.

<sup>c</sup> Hefei National Laboratory for Physical Sciences at the Microscale, University of Science and Technology of China, Hefei 230026, P. R. China.

<sup>d</sup> CAS Center for Excellence in Comparative Planetology, Hefei 230026, P. R. China.

Corresponding authors: Dr. Qi-Zhi Yao, Prof. Dr. Gen-Tao Zhou


Tel.: 86 551 63600533

Email: qzyao@ustc.edu.cn, gtzhou@ustc.edu.cn


### Preparation of boehmite template

In a typical procedure, 2344 mg of  $\text{Al}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$  was first dissolved into 50 mL of purified water, followed by the addition of 300 mg of  $(\text{NH}_2)_2\text{CO}$  under continuously stirring at room temperature to form a clear solution. The solution was then transferred to Teflon-lined stainless-steel autoclave (100 ml capacity). The hydrothermal vessel was heated to 175 °C, and maintained for 4 hours. After natural cooling, the product was collected by centrifugation, washed three times with purified water, and dried in a vacuum oven at 60 °C overnight.


The synthesized template boehmite was examined by XRD, FESEM and TEM techniques. As can be seen from Fig. S1a, the peaks at  $2\theta$  values of 14.4° (020), 28.1° (120), 38.3° (031), 45.7° (131), 48.9° (051), 49.2° (200), 51.5° (220), 55.2° (151), 60.5° (080), 64.0° (231), 64.9° (002), 66.1° (180) and 71.9° (251) match well with orthorhombic boehmite phase (JCPDS No. 21-1307). The FESEM and TEM images in Fig. S1b and c show that the synthesized boehmite owns a nanoplates-like structure, which are consistent with the results reported by Dubey et al. (2017). These results confirmed that the template boehmite was successfully synthesized.



**Fig. S1** XRD pattern (a), FESEM (b) and TEM (c) images of the synthesized boehmite template.



**Fig. S2** EDS spectrum of the synthesized product. Insets are corresponding analytical area and EDS datasheet, respectively.



**Fig. S3** XRD patterns of the products synthesized at 60 (a), 80 (b) and 150  $^{\circ}$ C (c) for different reaction times. Red asterisks (\*) represent  $\gamma$ -GaOOH and blue hearts (♥) represent  $\alpha$ -GaOOH

As shown in Fig. S3, the peaks corresponding to the characteristic peaks of  $\gamma$ -GaOOH marked by red asterisks appear in all temperature conditions, indicating that  $\gamma$ -GaOOH can be

obtained at the set temperature. In addition, with the increase of aging temperature, the occurrence time of the characteristic peaks of  $\gamma$ -GaOOH becomes shorter and shorter, from 12 days at 60 °C to 1 day and 1 hour at 80 and 150 °C, respectively, indicating that the  $\gamma$ -GaOOH is easier to form at higher temperature. Moreover, besides the diffraction peaks indexed to boehmite and  $\gamma$ -GaOOH, a small diffraction peak belonging to  $\alpha$ -GaOOH (JCPDS No. 06-0180) marked by blue heart in Fig. S3b and c appears, which indicate that higher temperature is not only conducive to the formation of  $\gamma$ -GaOOH, but also will leads to the formation of  $\alpha$ -GaOOH under the conditions explored in this work.

## References

Dubey, S.P., Dwivedi, A.D., Sillanpää, M., Lee, H., Kwon, Y.N., and Lee, C. (2017) Adsorption of As(V) by boehmite and alumina of different morphologies prepared under hydrothermal conditions. *Chemosphere*, 169, 99–106.