

Yangite, $\text{PbMnSi}_3\text{O}_8 \cdot \text{H}_2\text{O}$, a new mineral species with double wollastonite silicate chains, from the Kombat mine, Namibia

**ROBERT T. DOWNS¹, WILLIAM W. PINCH², RICHARD M. THOMPSON^{3,*}, STANLEY H. EVANS¹,
AND LAUREN MEGAW¹**

¹Department of Geosciences, University of Arizona, 1040 East 4th Street, Tucson, Arizona 85721, U.S.A.

²19 Stonebridge Lane, Pittsford, New York 14534, U.S.A.

³School of Information, University of Arizona, 1103 East 2nd Street, Tucson, Arizona 85721, U.S.A.

ABSTRACT

A new chain-silicate mineral species, yangite, ideally $\text{PbMnSi}_3\text{O}_8 \cdot \text{H}_2\text{O}$, has been found on a specimen from the Kombat mine, Otavi Valley, Namibia. Associated minerals are melanotekite and rhodochrosite. Yangite is colorless to pale brown in transmitted light, transparent with white streak and vitreous luster. Broken pieces of yangite crystals are bladed or platy, and elongated along [010]. It is sectile with a Mohs hardness of ~5; cleavage is perfect on {101} and no twinning or parting was observed. The measured and calculated densities are 4.14(3) and 4.16 g/cm³, respectively. Optically, yangite is biaxial (–), with $n_\alpha = 1.690(1)$, $n_\beta = 1.699(1)$, $n_\gamma = 1.705(1)$, $Y = b$, $Z \wedge c = 11^\circ$, and $2V_{\text{meas}} = 77(2)^\circ$. It is insoluble in water, acetone, and hydrochloric acid. An electron microprobe analysis demonstrated that the sample was relatively pure, yielding the empirical formula (with calculated H₂O) $\text{Pb}_{1.00}\text{Mn}_{1.00}^{2+}\text{Si}_{3.00}\text{O}_8 \cdot \text{H}_2\text{O}$. Yangite is triclinic and exhibits space group symmetry $P\bar{1}$ with unit-cell parameters $a = 9.6015(9)$, $b = 7.2712(7)$, $c = 7.9833(8)$ Å, $\alpha = 105.910(4)$, $\beta = 118.229(4)$, $\gamma = 109.935(5)^\circ$, and $V = 392.69(7)$ Å³. Its crystal structure is based on a skeleton of double wollastonite SiO₄ tetrahedral chains oriented parallel to [010] and interlinked with ribbons of Mn- and Pb-polyhedra. Yangite represents the first chain silicate with two-connected double chains and possesses all of the structural features of a hypothetical triclinic Ca₂Si₃O₈·2H₂O phase proposed by Merlini and Bonaccorsi (2008) as a derivative of the okenite structure. The difference in the H₂O component between the hypothetical phase and yangite likely is a consequence of the larger Pb²⁺ with its lone-pair electrons in yangite replacing the smaller Ca²⁺ in the hypothetical phase.

Keywords: Yangite, chain silicate, wollastonite chains, crystal structure, X-ray diffraction, Raman spectra