

## Elasticity of single-crystal periclase at high pressure and temperature: The effect of iron on the elasticity and seismic parameters of ferropericlase in the lower mantle

DAWEI FAN<sup>1,2,3,\*</sup>, SUYU FU<sup>2</sup>, JING YANG<sup>2,4</sup>, SERGEY N. TKACHEV<sup>5</sup>, VITALI B. PRAKAPENKA<sup>5</sup>, AND JUNG-FU LIN<sup>2,\*</sup>

<sup>1</sup>Key Laboratory of High Temperature and High Pressure Study of the Earth's Interior, Institute of Geochemistry, Chinese Academy of Sciences, Guiyang, Guizhou 550081, China ORCID 0000-0001-7840-2075

<sup>2</sup>Department of Geological Sciences, Jackson School of Geosciences, The University of Texas at Austin, Austin, Texas 78712, U.S.A.

<sup>3</sup>Center for High Pressure Science and Technology Advanced Research (HPSTAR), Shanghai 130012, China

<sup>4</sup>Geophysical Laboratory, Carnegie Institution of Washington, Washington, D.C. 20015, U.S.A.

<sup>5</sup>Center for Advanced Radiation Sources, University of Chicago, Chicago, Illinois 60437, U.S.A.

### ABSTRACT

In this study, we measured the sound velocities of single-crystal periclase by Brillouin light scattering (BLS) combined with in situ synchrotron X-ray diffraction (XRD) up to ~30 GPa and 900 K in an externally heated diamond-anvil cell (EHDAC). Our experimental results were used to evaluate the combined effects of pressure and temperature on the elastic moduli of single-crystal periclase using third-order Eulerian finite-strain equations. All of the elastic moduli increased with increasing pressure but decreased with increasing temperature, except the off-diagonal modulus  $C_{12}$ , which remained almost constant up to ~30 GPa and 900 K. The derived aggregate adiabatic bulk and shear moduli ( $K_{S0}$ ,  $G_0$ ) at ambient conditions were 162.8( $\pm$ 0.2) and 130.3( $\pm$ 0.2) GPa, respectively, consistent with literature results. The pressure derivatives of the bulk [ $(\partial K_S/\partial P)_{300\text{ K}}$ ] and shear moduli [ $(\partial G/\partial P)_{300\text{ K}}$ ] at ambient conditions were 3.94( $\pm$ 0.05) and 2.17( $\pm$ 0.02), respectively, whereas the temperature derivatives of these moduli [ $(\partial K_S/\partial T)_P$ ] and [ $(\partial G/\partial T)_P$ ] at ambient conditions were  $-0.025(\pm 0.001)$  and  $-0.020(\pm 0.001)$  GPa/K, respectively. A comparison of our experimental results with the high-pressure ( $P$ ) and high-temperature ( $T$ ) elastic moduli of ferropericlase (Fp) in the literature showed that all the elastic moduli of Fp were linearly correlated with the FeO content up to approximately 20 mol%. These results allowed us to build a comprehensive thermoelastic model for Fp to evaluate the effect of Fe-Mg substitution on the elasticity and seismic parameters of Fp at the relevant  $P$ - $T$  conditions of the lower mantle. Our modeling results showed that both the increase of the Fe content in Fp and the increasing depth could change the compressional wave anisotropy ( $AV_p$ ) and shear wave splitting anisotropy ( $AV_s$ ) of Fp in the upper parts of the lower mantle. Furthermore, using our modeling results here, we also evaluated the contribution of Fp to seismic lateral heterogeneities of thermal or chemical origin in the lower mantle. Both the thermally induced and Fe-induced heterogeneities ratios ( $R_{S/P} = \partial \ln V_s / \partial \ln V_p$ ) of Fp from 670 to 1250 km along a representative lower mantle geotherm increased by ~2–5% and ~15%, respectively. The thermally induced  $R_{S/P}$  value of Fp20 is ~30% higher than Fp10, indicating that the Fe content has a significant effect on the thermally induced  $R_{S/P}$  of Fp. Compared to the seismic observation results ( $R_{S/P} = 1.7$ – $2.0$ ) in the upper regions of the lower mantle, the Fe-induced  $R_{S/P}$  value of Fp is more compatible than the thermally induced  $R_{S/P}$  value of Fp20 (the expected composition of Fp in the lower mantle) within their uncertainties. Thus, we propose that Fe-induced lateral heterogeneities can significantly contribute to the observed seismic lateral heterogeneities in the Earth's lower mantle (670–1250 km).

**Keywords:** Elasticity, periclase, Brillouin light scattering, lower mantle, diamond anvil cell