

## **Crystal chemistry of Cr-spinels from the lherzolite mantle peridotite of Ronda (Spain)**

**DAVIDE LENAZ,<sup>1,\*</sup> ANGELO DE MIN,<sup>1</sup> GIORGIO GARUTI,<sup>2</sup> FEDERICA ZACCARINI,<sup>2</sup>  
AND FRANCESCO PRINCIVALLE<sup>1</sup>**

<sup>1</sup>Department of Geosciences, University of Trieste, Via Weiss 8, 34127 Trieste, Italy

<sup>2</sup>Department of Applied Geosciences and Geophysics, University of Leoben, Peter Tunner Strasse 5, 8700 Leoben, Austria

### **ABSTRACT**

The crystal chemistry of some Cr-spinels from the lherzolite body of the Ronda peridotite in southern Spain has been investigated. Cell edge spans between 8.1692(2) and 8.2367(1) Å, while the oxygen positional parameter  $u$  ranges between 0.26306(7) and 0.26351(7). By using the Princivalle thermometer (1999), an intracrystalline closure temperature between 640 and 840 °C has been calculated. The higher temperatures are very close to the intercrystalline temperatures based on the olivine-spinel thermometer calculated by Woodland et al. (2006) for the Ronda orogenic lherzolites suggesting that the intracrystalline closure occurred soon after the intercrystalline closure. By comparison with Cr-spinels from lherzolite mantle xenoliths, it should be noted that: (1) the oxygen positional parameter can be linearly related to the intracrystalline temperature for both mantle peridotite and mantle xenolith Cr-spinels; (2) the intracrystalline closure temperature is reached faster and is higher in Cr-spinels in mantle xenoliths; and (3) Cr content is linearly related to  $u$  in mantle peridotite, but not in mantle xenoliths, suggesting the  $u$  value in Cr-spinels from mantle peridotite is driven solely by the chemistry of the spinels.

**Keywords:** Cr-spinel, crystal chemistry, intracrystalline closure temperature, mantle peridotite, Ronda (Spain)