

## **In-situ mid/far micro-FTIR spectroscopy to trace pressure-induced phase transitions in strontium feldspar and wadsleyite**

**MARIA MROSKO,<sup>1,\*</sup> MONIKA KOCH-MÜLLER,<sup>1</sup> AND ULRICH SCHADE<sup>2</sup>**

<sup>1</sup>Helmholtz-Zentrum Potsdam, Deutsches GeoForschungsZentrum-GFZ, Telegrafenberg, 14473 Potsdam, Germany

<sup>2</sup>Helmholtz-Zentrum Berlin für Materialien und Energie, Albert-Einstein-Strasse 15, 12489 Berlin, Germany

### **ABSTRACT**

As representatives of nominally anhydrous minerals (NAMs) in the crust and mantle the pressure-dependent behavior of strontium feldspar and wadsleyite, containing different amounts of water, was studied in a diamond-anvil cell via mid and far IR spectroscopy up to 24 GPa. The samples were synthesized in a piston-cylinder press at 2 GPa/700 °C (strontium feldspar) and in a multi-anvil apparatus at 13.8 GPa/1000 °C (dry wadsleyite) and 13.2 GPa/1150 °C (hydrous wadsleyite). The water content of the samples was determined by polarized FTIR and Raman spectroscopy. The strontium feldspar crystals (up to 300 µm) contained about 1100(100) wt ppm water. The hydrous wadsleyite crystals (up to 240 µm) contained 12 500(900) wt ppm water. The synthesis of dry wadsleyite yielded a fine-grained powder. A new THz/FIR-microscope for the synchrotron source BESSY was developed to conduct the diamond-anvil cell measurements in the far IR region. Conventional in-situ mid IR spectroscopy was also performed on all samples. The measurements on strontium feldspar showed a phase transformation at 6.5(5) GPa (space group  $I2/c$  to  $P2_1/c$ ). The wadsleyite analyses revealed a phase transition at approximately 8.4(7) GPa in the hydrous and approximately 10.0(7) GPa in the dry sample. It probably represents a transition from an orthorhombic to a monoclinic structure. The high amount of water incorporated in the hydrous wadsleyite shifts the transformation toward lower pressures compared to the dry one. By comparison, the relatively low amount of water in strontium feldspar does not change the stability relations compared to the dry one. Therefore, water incorporation in nominally anhydrous minerals may have an effect on the pressure of phase transitions, whereas the extent of that influence strongly depends on the structure of the phase and the amount of water carried within the mineral.

**Keywords:** Wadsleyite, strontium feldspar, mid/far FTIR spectroscopy, phase transition, water