

Supporting information for:

## **Effect of iron and trivalent cations on OH-defects in olivine**

Marc Blanchard<sup>1</sup>, Jannick Ingrin<sup>2</sup>, Etienne Balan<sup>1</sup>, István Kovács<sup>3</sup>, Anthony C. Withers<sup>4</sup>


<sup>1</sup> Institut de Minéralogie, de Physique des Matériaux, et de Cosmochimie (IMPMC), Sorbonne Universités, UPMC Univ Paris 06, UMR CNRS 7590, Muséum National d'Histoire Naturelle, UMR IRD 206, 4 place Jussieu, F-75005 Paris, France

<sup>2</sup> UMET, UMR CNRS 8207, Université de Lille1, Bât. C6, 59655 Villeneuve d'Ascq, France


<sup>3</sup> Geological and Geophysical Institute of Hungary, Budapest, Stefánia street 14, H-1143, Hungary

<sup>4</sup> Department of Earth Sciences, University of Western Ontario, London, ON, Canada N6A 5B7


\*Communicating Author: Marc Blanchard. E-mail: marc.blanchard@impmc.upmc.fr




**FIGURE S1.** Theoretical polarized infrared spectra of  $(2H)_M^x$  defects.  $(2H)_{M1}^x$  corresponds to the M1 configuration in Balan et al. (2011).



**FIGURE S2.** Structural models of  $(2H)_{M2}^x$  defects, with Mg in yellow, Si in blue, O in red and H in white (color online).



**FIGURE S3.** Theoretical polarized infrared spectra of  $(\text{Fe}+1\text{H})_{\text{Si}}^{\times}$  defect (left) and the corresponding structural model (right).



**FIGURE S4.** Theoretical polarized infrared spectra of  $[(\text{Fe})_{\text{M1}}^{\bullet}(\text{3H})'_{\text{Si}}]^{\times}$  defect (left) and the corresponding structural model (right).