

Immiscible-melt inclusions in corundum megacrysts: Microanalyses and geological implications

**XI-SHENG XU^{1,2,*}, XIAO-MING CHEN¹, WILLIAM L. GRIFFIN^{1,2,†}, SUZANNE Y. O'REILLY^{1,2},
XI-SONG ZHANG¹, AND LI-HUI CHEN¹**

¹State Key Laboratory for Mineral Deposits Research, School of Earth Sciences and Engineering, Nanjing University, Nanjing 210023, China

²ARC Centre of Excellence for Core to Crust Fluid Systems and GEMOC, Department Earth and Environmental Sciences, Macquarie University, New South Wales 2109, Australia

ABSTRACT

Controversies on the origin of zircon, corundum, titanomagnetite, and quartz megacrysts in alkali basalts mostly reflect the lack of direct evidence of a “melt reservoir” required for their formation. Various mineral megacrysts are carried up by Cenozoic (mostly younger than 25 Ma) alkali basalts that extend more than 4000 km along eastern China. Here we report unusual inclusions in corundum megacrysts from Changle, and we attribute their origin to the existence of a FeO*-SiO₂-Al₂O₃-ZrO₂-rich melt. The inclusions, analyzed using electron microprobe and Raman microscopy, may be divided into two types. Type I inclusions are dominated by glassy materials, may exhibit a dark part in backscattered electron (BSE) images composed of quartz, corundum, and an amorphous substance (AS-1), and a bright part in BSE images composed of baddeleyite and a second distinct amorphous substance (AS-2). Compared with AS-1, AS-2 has higher concentrations of ZrO₂ and FeO* but lower concentrations of Al₂O₃ and SiO₂. We argue that the formation temperature of Type I inclusions is ~1200 °C, and the generation of their bright and dark parts in BSE images may be attributed to the coexistence of immiscible melts. Type II inclusions are composed of zircon, quartz, and an amorphous substance (AS-3). Both types of inclusions might be derived from a similar parent melt, which is FeO*-SiO₂-Al₂O₃-ZrO₂-rich. New secondary ion mass spectroscopy (SIMS) in situ U-Pb ages of 18 Ma and 13–14 Ma for zircon inclusions suggest that the corundum megacrysts, occurring in basaltic host rocks distributed along the middle segment of the north and south-trending Tanlu fault zone, formed from precursor residual magmas related to underplating basalts stalled at the crust-mantle boundary, and were brought to the surface by entrainment in later basalts. This study provides new insights into the genesis of the corundum-related megacryst suite.

Keywords: Corundum megacrysts, immiscible-melt inclusions, underplating basalts, geochemical analysis, BSE imaging