

**PARTIAL MELTING AND P-T EVOLUTION OF ECLOGITE-FACIES
METAPELITIC MIGMATITES FROM THE EGERE TERRANE (CENTRAL
HOGGAR, SOUTH ALGERIA)**

**Amar Arab, Gaston Godard, Khadidja Ouzegane, Antonio Acosta-Vigil, Jean-Robert
Kiénast, Manuel J. Román-Alpiste, Carlos J. Garrido, Amar Drareni**

SUPPLEMENTARY MATERIALS

1 – METHODS

2 – FIGURE

Figure A

3 – ANALYZES

Table A

Table B

Table C

Table D

Table E

Table F

1 – METHODS

A selection of high-grade metapelite samples was studied under scanning electron microscope at Paris-Sorbonne University (<http://meb.ufr918.upmc.fr>), producing in particular backscattered electron images and X-ray elemental maps. Chemical analyses of minerals (Tables B to G) were undertaken using a CAMECA SX100 electron microprobe at Camparis laboratory in Paris (<http://camparis.ecceterra.fr/>) with operating conditions of 15 kV and 10 nA; natural standards were used for calibration. Whole-rock analyses of major elements (Table A) were performed by X-ray fluorescence (XRF-M01) at the C.R.P.G. service (Nancy, France; see <http://www.crpq.cnrs-nancy.fr/SARM> for the analytical protocols).

Crystallized melt inclusions (“nanogranitoids”) and other melt-related inclusions in garnet were characterized on thin sections using conventional microscope petrography and Zeiss MA-15 and Quanta 400 SEMs at the Instituto Andaluz de Ciencias de la Tierra (CSIC-Universidad de Granada) and Centro de Instrumentación Científica (Universidad de Granada).

P – T pseudosections were calculated in the system MnO – Na_2O – CaO – K_2O – FeO – MgO – Al_2O_3 – SiO_2 – H_2O – TiO_2 – O , which is a near comprehensive compositional analog of natural metapelitic rocks (White et al. 2014a), using the thermodynamic software THERMOCALC v3.40 (Powell and Holland 1988) and an updated version of the Holland and Powell (2011) data set (file tc-ds62.txt). Fe^{3+} in the bulk compositions is fixed at 10 mol.% of total iron, as advised by Diener and Powell (2010). The activity-composition models used are those of White et al. (2014a, b). Quartz, kyanite, sillimanite, rutile and fluid (H_2O) are pure phases. Isopleth curves for selected minerals show the evolution of their chemical composition with respect to P and T (Fig. 7). The following notations are used: $x(\text{Grt}) = \text{Fe}/(\text{Fe} + \text{Mg})$; $z(\text{Grt}) = \text{Ca}/(\text{Ca} + \text{Fe} + \text{Mg} + \text{Mn})$; $x(\text{Bt}) = \text{Fe}/(\text{Fe} + \text{Mg})$; $x(\text{St}) = \text{Fe}/(\text{Fe} + \text{Mg})$; $\text{Si}(\text{Ph}) = \text{Si}$.

The mineral abbreviations used in the text are from Whitney and Evans (2010).

Bibliography:

- Diener, J.F.A., and Powell, R. (2010) Influence of ferric iron on the stability of mineral assemblages. *Journal of Metamorphic Geology*, 28(6), 599-613.
- Holland, T., and Powell, R. (2011) An improved and extended internally consistent thermodynamic dataset for phases of petrological interest, involving a new equation of state for solids. *Journal of Metamorphic Geology*, 29(3), 333-383.
- Powell, R., and Holland, T.J.B. (1988) An internally consistent dataset with uncertainties and correlations: 3. Applications to geobarometry, worked examples and a computer program. *Journal of Metamorphic Geology*, 6(2), 173-204.
- White, R., Powell, R., Holland, T., Johnson, T., and Green, E. (2014a) New mineral activity–composition relations for thermodynamic calculations in metapelitic systems. *Journal of Metamorphic Geology*, 32(3), 261-286.
- White, R., Powell, R., and Johnson, T. (2014b) The effect of Mn on mineral stability in metapelites revisited: New a – x relations for manganese-bearing minerals. *Journal of Metamorphic Geology*, 32(8), 809-828.
- Whitney, D.L., and Evans, B.W. (2010) Abbreviations for names of rock-forming minerals. *American Mineralogist*, 95(1), 185-187.

2 – FIGURE




Figure A. Photomicrographs of the rocks studied.

Scale bars correspond to a length of 5 mm; thin sections are seen in plane-polarized (left) and cross-polarized (right) lights. **(a, b)** Phengite-rich metapelite. **(c, d)** Biotite-rich metapelite, where biotite mainly developed after phengite. **(e, f)** Phengite-rich metapelite. **(g, h)** Metapelite rich in kyanite (yellow arrows in g) and phengite. The samples show leucocratic bands (*i.e.*, leucosomes) alternating with darker bands rich in garnet and micas. The foliation visible in the matrix, mainly defined by quartz and phengite, wraps around garnet porphyroblasts. The inclusions in garnet define an internal foliation, either sigmoidal (red arrow in b) or straight (green arrows in c, e and g), at variable angles with respect to the foliation in the matrix (see g and f). In many areas, biotite-rich post-foliation fine-grained M_3 assemblages replace the garnet porphyroblasts (blue arrows in h) and especially the phengite flakes Ph_2 (c and d).

3- ANALYZES

Table A: Studied samples coordinates and bulk-rock compositions (in mol %) used for modeling the pseudosections

Sample	SiO ₂	TiO ₂	Al ₂ O ₃	FeO	MgO	MnO	CaO	Na ₂ O	K ₂ O	O	H ₂ O	Longitude	Latitude
T52	68.18	1.20	10.80	9.95	1.29	0.11	2.67	1.37	2.14	0.50	1.73	5° 51' 34.65" E	25° 58' 31.93" N
T78	60.31	0.99	11.08	6.87	1.71	0.08	1.63	1.31	2.47	0.43	13.12	5° 51' 29.41" E	25° 58' 23.47" N
T133	70.35	1.07	10.65	8.04	2.48	0.01	2.70	1.04	2.12	0.50	1.06	5° 51' 35.56" E	25° 58' 49.78" N
T138	73.34	1.15	8.13	9.03	1.13	0.05	2.18	1.02	2.06	0.50	1.36	5° 51' 35.40" E	25° 58' 42.80" N
T51	—	—	—	—	—	—	—	—	—	—	—	5° 51' 43.11" E	25° 58' 32.17" N
T77	—	—	—	—	—	—	—	—	—	—	—	5° 51' 29.41" E	25° 58' 27.84" N
T100	—	—	—	—	—	—	—	—	—	—	—	5° 51' 35.60" E	25° 58' 41.60" N
T109	—	—	—	—	—	—	—	—	—	—	—	5° 51' 35.40" E	25° 58' 43.50" N

Table B: Representative microprobe analyzes of garnet

Cationic formulas are calculated on the basis of 12 O

Sample	T52	T77	T133		T138				T78	T138	
	Grt ₁ (core)				Grt (rim)		Grt (outer rim)		Grt ₃		
SiO ₂	37.68	37.94	37.22	37.35	37.59	37.85	37.71	36.35	37.70	37.06	37.31
TiO ₂	0.00	0.07	0.10	0.11	0.03	0.04	0.01	0.02	0.00	0.00	0.00
Al ₂ O ₃	21.09	20.58	21.22	20.83	20.85	20.99	21.04	20.06	21.09	20.88	20.93
Cr ₂ O ₃	0.02	0.00	0.00	0.04	0.00	0.01	0.00	0.02	0.03	0.02	0.01
FeO _t	32.79	30.34	30.81	29.94	34.69	34.00	35.03	32.86	35.87	38.19	38.17
MnO	0.19	0.76	2.06	2.31	0.81	0.81	0.56	0.85	1.12	0.14	0.12
MgO	2.65	2.09	0.89	0.95	2.69	2.51	2.50	2.49	3.49	1.61	1.67
CaO	6.92	8.74	8.77	8.95	3.34	4.43	4.83	5.00	1.91	3.06	3.20
Na ₂ O	0.08	0.06	0.00	0.00	0.00	0.00	0.01	0.03	0.03	0.00	0.02
K ₂ O	0.02	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01
Total	101.44	100.71	101.08	100.48	100.01	100.68	101.76	97.80	101.31	100.98	101.47
Si	2.980	3.017	2.971	2.993	3.020	3.018	2.991	2.998	2.997	2.989	2.993
Al	1.966	1.929	1.996	1.967	1.974	1.973	1.966	1.950	1.976	1.985	1.979
Cr	0.000	0.000	0.006	0.007	0.002	0.002	0.001	0.001	0.002	0.001	0.001
Ti	0.001	0.004	0.000	0.003	0.000	0.001	0.000	0.001	0.000	0.000	0.000
Fe ³⁺	0.033	0.067	0.000	0.023	0.024	0.024	0.033	0.048	0.022	0.014	0.021
Fe ²⁺	2.136	1.950	2.057	1.983	2.306	2.243	2.290	2.218	2.363	2.562	2.540
Mn	0.013	0.051	0.139	0.157	0.055	0.055	0.038	0.059	0.076	0.010	0.008
Mg	0.312	0.248	0.106	0.113	0.322	0.298	0.296	0.306	0.414	0.194	0.200
Ca	0.586	0.745	0.750	0.768	0.287	0.378	0.410	0.442	0.164	0.264	0.275
Na	0.012	0.009	0.000	0.000	0.000	0.000	0.002	0.005	0.005	0.000	0.003
K	0.002	0.000	0.001	0.000	0.001	0.000	0.001	0.000	0.001	0.000	0.001
Σ Cat.	8.043	8.020	8.026	8.015	7.992	7.993	8.027	8.028	8.017	8.018	8.020
X _{Fe}	0.868	0.867	0.893	0.880	0.859	0.864	0.873	0.859	0.829	0.927	0.924
X _{Alm}	0.701	0.651	0.674	0.656	0.776	0.754	0.755	0.733	0.784	0.846	0.840
X _{Sps}	0.004	0.017	0.046	0.052	0.019	0.018	0.012	0.020	0.025	0.003	0.003
X _{Prp}	0.103	0.083	0.035	0.038	0.108	0.100	0.097	0.101	0.137	0.064	0.066
X _{Grs}	0.175	0.215	0.246	0.241	0.085	0.115	0.119	0.122	0.042	0.080	0.080
X _{Adr}	0.016	0.034	0.000	0.012	0.012	0.012	0.016	0.024	0.011	0.007	0.010

Table C: Representative microprobe analyzes of white micas

Cationic formulas are calculated on the basis of $O_{20}(OH,F,Cl)_4$ * Ms_3 within the Bt_3-Pl_3 symplectites after Ph_2 ;** Estimated composition of Ti-rich phengite before Ilm_3 exsolution, reconstructed by scanning with the microprobe (see text);*** H_2O calculated by stoichiometry; O equivalent to F+Cl subtracted from the total weight.

Sample	T138				T52		T138			T52		T138
	Ph ₂ core		Ph ₂ rim		Ph ₁ in Grt ₁		Ms ₃ after Ph ₂ *		Ms ₃ matrix		Ti-Ph **	
SiO ₂	47.79	47.67	43.68	46.34	46.81	46.76	43.68	45.30	45.68	45.21	47.47	38.04
TiO ₂	1.22	1.23	0.92	1.54	1.46	1.77	0.92	0.97	0.39	1.28	1.44	5.88
Al ₂ O ₃	31.86	31.96	35.85	32.98	33.24	33.70	35.85	36.18	36.94	34.33	34.29	29.28
Cr ₂ O ₃	0.05	0.01	0.01	0.03	0.05	0.06	0.01	0.01	0.00	0.04	0.06	0.04
FeO	2.09	1.90	1.33	1.71	1.44	1.49	1.33	1.46	1.47	1.05	1.32	5.25
MnO	0.00	0.04	0.00	0.04	0.00	0.00	0.00	0.02	0.01	0.02	0.02	0.04
MgO	1.38	1.24	0.29	0.86	0.90	0.91	0.29	0.29	0.30	0.61	0.80	0.22
CaO	0.01	0.02	0.15	0.11	0.01	0.01	0.15	0.07	0.02	0.01	0.01	0.14
Na ₂ O	0.46	0.37	0.47	0.40	0.51	0.47	0.47	0.46	0.69	0.53	0.54	0.48
K ₂ O	11.10	11.36	11.03	11.31	10.95	11.09	11.03	11.98	10.62	11.17	11.12	8.50
F	0.00	0.00	0.00	0.00	0.00	0.40	0.00	0.00	0.00	0.00	0.00	0.00
Cl	0.00	0.00	0.00	0.00	0.02	0.01	0.00	0.00	0.02	0.01	0.00	0.00
H ₂ O***	4.50	4.49	4.40	4.46	4.48	4.34	4.40	4.52	4.53	4.57	4.43	4.02
Total	100.46	100.29	98.13	99.78	99.88	100.84	98.13	101.26	100.67	98.83	101.50	91.89
Si	6.367	6.364	5.956	6.224	6.253	6.195	5.956	6.008	6.038	6.119	6.226	5.672
Al ^{IV}	1.633	1.636	2.043	1.775	1.746	1.804	2.044	1.992	1.961	1.880	1.773	2.328
Al ^{VI}	3.369	3.393	3.719	3.444	3.487	3.457	3.718	3.664	3.795	3.596	3.526	2.819
Cr	0.005	0.002	0.001	0.003	0.005	0.007	0.001	0.001	0.000	0.004	0.005	0.005
Ti	0.122	0.123	0.094	0.155	0.146	0.176	0.094	0.097	0.038	0.130	0.142	0.660
Fe	0.233	0.212	0.151	0.191	0.161	0.165	0.152	0.162	0.163	0.119	0.145	0.655
Mn	0.000	0.005	0.000	0.004	0.000	0.000	0.000	0.002	0.001	0.002	0.002	0.005
Mg	0.274	0.247	0.059	0.172	0.178	0.179	0.059	0.057	0.058	0.122	0.155	0.050
Ca	0.001	0.002	0.021	0.015	0.001	0.001	0.022	0.010	0.002	0.001	0.001	0.022
Na	0.119	0.095	0.123	0.104	0.132	0.119	0.124	0.118	0.177	0.139	0.137	0.138
K	1.886	1.934	1.918	1.937	1.865	1.874	1.919	2.027	1.791	1.928	1.860	1.618
Σ cat.	14.010	14.013	14.089	14.030	13.979	13.988	14.089	14.139	14.028	14.044	13.978	13.970
X _{Fe}	0.46	0.46	0.72	0.53	0.47	0.48	0.72	0.73	0.74	0.49	0.48	0.92

Table D: Representative microprobe analyzes of biotite

Cationic formulas are calculated on the basis of $O_{20}(OH,F,Cl)_4$.* H_2O calculated by stoichiometry; O equivalent to F+Cl subtracted from the total weight.

Sample	T133		T77	T52	T138				
	Bt ₃ , after Grt, with Pl ₃ (partially chloritized)			Bt ₃ , after Ph ₂ and in the matrix, with Pl ₃ ±St ₃ ±Sil ₃	Bt ₃ , as exsolution lamellas in Ph ₂				
SiO ₂	36.11	36.19	34.64	34.07	33.26	33.07	33.49	34.17	34.48
TiO ₂	3.10	3.02	2.97	3.42	4.05	3.03	2.27	2.90	2.98
Al ₂ O ₃	19.45	20.36	18.48	19.15	19.06	19.28	19.78	19.45	20.23
Cr ₂ O ₃	0.01	0.03	0.03	0.14	0.05	0.15	0.12	0.06	0.10
FeO _t	15.19	15.38	20.71	22.15	26.85	29.74	24.10	23.44	23.15
MnO	0.05	0.08	0.05	0.04	0.00	0.00	0.00	0.03	0.01
MgO	11.23	10.86	8.53	6.17	3.85	2.58	6.74	6.52	5.79
CaO	0.03	0.00	0.02	0.02	0.06	0.00	0.27	0.01	0.08
Na ₂ O	0.17	0.11	0.18	0.19	0.20	0.17	0.18	0.17	0.17
K ₂ O	9.74	10.21	9.62	10.24	9.45	9.43	9.14	10.25	10.22
F	0.00	0.00	0.32	0.05	0.00	0.00	0.00	0.53	1.14
Cl	0.02	0.00	0.00	0.01	0.01	0.02	0.00	0.00	0.00
H ₂ O*	3.93	4.05	3.74	3.84	3.83	3.80	3.87	3.65	3.39
Total	99.10	100.29	99.16	99.49	100.67	101.27	99.96	100.96	101.26
Si	5.398	5.354	5.334	5.286	5.195	5.201	5.188	5.249	5.264
Al ^{IV}	2.602	2.645	2.666	2.713	2.805	2.798	2.812	2.750	2.735
Al ^{VI}	0.824	0.903	0.688	0.786	0.704	0.776	0.801	0.770	0.905
Cr	0.001	0.004	0.004	0.017	0.006	0.018	0.014	0.006	0.012
Ti	0.349	0.336	0.344	0.399	0.475	0.358	0.265	0.334	0.341
Fe	1.899	1.903	2.667	2.873	3.507	3.911	3.122	3.011	2.955
Mn	0.006	0.010	0.007	0.004	0.000	0.000	0.000	0.003	0.001
Mg	2.502	2.395	1.958	1.427	0.897	0.605	1.556	1.492	1.316
Ca	0.005	0.000	0.003	0.003	0.010	0.000	0.045	0.001	0.013
Na	0.049	0.032	0.054	0.057	0.060	0.051	0.055	0.050	0.050
K	1.857	1.927	1.890	2.027	1.884	1.892	1.806	2.008	1.989
Σ cat.	15.493	15.510	15.615	15.600	15.544	15.620	15.664	15.680	15.590
X _{Fe}	0.431	0.44	0.58	0.67	0.80	0.87	0.67	0.67	0.69

Table E: Representative microprobe analyzes of feldspars.

Cationic formulas are calculated on the basis of 8 O. Fe is considered as Fe³⁺.

Sample	T138								T52	T77	T78	T138	
	Pl ₃ , in pseudo-morph after Ph ₂		Pl, around Qz ₁ in Grt ₁	Pl, around Bt ₁ inclusion in Grt ₁		Pl, included in Grt ₁		Pl ₃ , associated with Bt ₃ ±St ₃ ±Kfs ₃ ±Sil ₃				Kfs ₃ , as corona between Ph ₂ & Qz ₂	
SiO₂	58.74	58.35	61.27	61.26	62.72	63.09	59.11	64.64	60.34	60.59	64.59	64.42	
TiO₂	0.01	0.07	0.01	0.01	0.00	0.03	0.00	0.00	0.05	0.02	0.00	0.00	
Al₂O₃	24.74	26.88	20.77	25.63	25.85	24.49	24.38	22.36	25.84	25.38	18.81	18.54	
Cr₂O₃	0.00	0.02	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.01	0.02	0.02	
Fe₂O₃t	0.32	0.59	0.68	0.48	0.43	0.68	0.33	0.11	0.01	0.16	0.06	0.03	
MnO	0.01	0.00	0.01	0.03	0.01	0.00	0.04	0.03	0.01	0.04	0.00	0.00	
MgO	0.00	0.10	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.01	
CaO	7.06	7.03	3.08	6.51	6.47	4.94	6.59	3.07	7.10	7.09	0.03	0.03	
Na₂O	8.60	7.29	10.30	7.17	6.17	9.03	8.28	10.27	8.00	8.03	0.56	0.46	
K₂O	0.13	1.02	0.12	0.14	0.19	0.20	0.19	0.28	0.07	0.13	16.10	16.88	
Total	102.21	101.35	96.26	101.17	101.84	102.46	98.95	100.76	101.42	101.45	100.19	100.39	
Si	2.647	2.587	2.826	2.689	2.717	2.736	2.672	2.835	2.654	2.667	2.980	2.980	
Al	1.314	1.405	1.129	1.326	1.320	1.252	1.299	1.156	1.340	1.317	1.023	1.011	
Fe³⁺	0.012	0.020	0.024	0.016	0.014	0.022	0.011	0.004	0.000	0.006	0.002	0.001	
Cr	0.000	0.001	0.000	0.000	0.000	0.000	0.001	0.000	0.000	0.000	0.001	0.001	
Ti	0.000	0.002	0.000	0.000	0.000	0.001	0.000	0.000	0.002	0.001	0.000	0.000	
Mn	0.000	0.000	0.000	0.001	0.000	0.000	0.002	0.001	0.000	0.002	0.000	0.000	
Mg	0.000	0.007	0.001	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.001	
Ca	0.341	0.334	0.152	0.306	0.300	0.230	0.319	0.144	0.335	0.335	0.001	0.001	
Na	0.752	0.627	0.921	0.610	0.518	0.759	0.726	0.873	0.682	0.686	0.050	0.041	
K	0.008	0.058	0.007	0.008	0.011	0.011	0.011	0.016	0.004	0.008	0.948	0.996	
Σ cat.	5.074	5.040	5.061	4.956	4.880	5.011	5.041	5.029	5.017	5.022	5.006	5.032	
X_{An}	0.312	0.328	0.141	0.331	0.362	0.230	0.302	0.140	0.328	0.326	0.001	0.001	
X_{Ab}	0.683	0.615	0.853	0.660	0.625	0.759	0.687	0.845	0.668	0.667	0.050	0.040	
X_{Or}	0.007	0.057	0.007	0.008	0.013	0.011	0.010	0.015	0.004	0.007	0.948	0.959	

Table F: Representative microprobe analyzes of staurolite.

Cationic formulas are calculated on the basis of $O_{46}(OH,F,Cl)_2$.* H_2O calculated by stoichiometry; O equivalent to F+Cl subtracted from the total weight.

Sample	T78				T138			T52	
	St ₁ (inclusion in Grt ₁)				St ₃ (matrix)				
SiO ₂	26.37	26.48	27.37	26.72	25.56	25.99	25.01	23.82	25.23
TiO ₂	0.71	0.87	0.72	0.64	0.63	0.52	0.56	0.69	0.49
Al ₂ O ₃	54.89	54.55	54.45	54.55	56.89	56.42	55.74	58.01	56.64
FeO	14.23	14.27	13.79	13.10	14.82	14.52	14.16	13.71	14.17
MnO	0.16	0.20	0.23	0.22	0.05	0.00	0.03	0.06	0.00
MgO	1.22	1.22	1.36	1.25	0.53	0.64	0.65	0.73	0.88
ZnO	0.89	0.93	1.38	1.09	1.03	1.16	0.88	0.61	0.62
CaO	0.06	0.07	0.02	0.04	0.02	0.01	0.02	0.04	0.00
Na ₂ O	0.08	0.13	0.11	0.10	0.05	0.06	0.04	0.10	0.12
K ₂ O	0.01	0.01	0.00	0.02	0.02	0.03	0.01	0.01	0.00
F	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00
Cl	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
H ₂ O*	1.08	1.06	1.07	1.04	1.06	1.06	1.04	1.05	1.05
Total	99.86	99.98	100.72	99.01	100.71	100.42	98.17	98.89	99.21
Si	7.329	7.519	7.704	7.620	7.206	7.334	7.214	6.808	7.184
Al	17.980	18.256	18.062	18.335	18.903	18.765	18.948	19.542	19.007
Ti	0.148	0.186	0.152	0.137	0.134	0.110	0.122	0.148	0.105
Fe	3.308	3.389	3.246	3.124	3.494	3.427	3.416	3.277	3.374
Mn	0.038	0.048	0.055	0.053	0.012	0.000	0.007	0.015	0.000
Mg	0.505	0.516	0.571	0.531	0.223	0.269	0.279	0.311	0.374
Zn	0.183	0.194	0.287	0.230	0.214	0.242	0.187	0.129	0.131
Ca	0.018	0.021	0.006	0.012	0.006	0.003	0.006	0.012	0.000
Na	0.043	0.072	0.060	0.055	0.027	0.033	0.022	0.055	0.066
K	0.004	0.004	0.000	0.007	0.007	0.011	0.004	0.004	0.000
Σ Cat.	29.556	30.205	30.143	30.106	30.226	30.195	30.204	30.302	30.241
X _{Mg}	0.13	0.12	0.14	0.13	0.06	0.07	0.07	0.08	0.10