

First-principles calculation of the infrared spectrum of hematite

MARC BLANCHARD,^{1,2,*} MICHELE LAZZERI,² FRANCESCO MAURI,² AND ETIENNE BALAN^{1,2}

¹IRD, 209 rue La Fayette, 75480 Paris Cedex 10, France

²Institut de Minéralogie et de Physique des Milieux Condensés (IMPMC), UMR CNRS 7590, Universités Paris VI et VII, IPGP, Campus Boucicaut, 140 rue de Lourmel, 75015 Paris, France

ABSTRACT

The theoretical infrared spectrum of hematite (α -Fe₂O₃) was computed using ab initio quantum mechanical calculations. Frequencies of the normal vibrational modes and Born effective charges were computed using the density functional theory (DFT) with and without the addition of a Hubbard U correction. The infrared reflection spectra of a single crystal of hematite were calculated as well as the infrared powder absorption spectrum using an electrostatic model that takes into account the shape of hematite particles. The theoretical behavior of the absorption bands is in agreement with experimental observations and provides a firm basis for the interpretation of the bands in terms of vibrational modes. Overall, results suggest that the use of DFT + U , which is necessary to describe correctly the electronic and magnetic properties of hematite, does not improve noticeably the prediction of vibrational properties.

Keywords: Iron oxide, density functional theory, infrared spectroscopy