

LETTER

Heat capacity and entropy of melanophlogite: Molecule-containing porosils in nature

CHARLES A. GEIGER,^{1,*} EDGAR DACHS,² AND MARIKO NAGASHIMA¹

¹Institut für Geowissenschaften, Abteilung Mineralogie, Christian-Albrechts-Universität zu Kiel, D-24098 Kiel, Germany

²Fachbereich Materialforschung und Physik, Abteilung Mineralogie, Universität Salzburg, Hellbrunnerstrasse 34, A-5020 Salzburg, Austria

ABSTRACT

The heat capacities of two different molecule-containing melanophlogites of approximate composition $46\text{SiO}_2 \cdot 1.80\text{CH}_4 \cdot 3.54\text{N}_2 \cdot 1.02\text{CO}_2$ from Mt. Hamilton, California, and $46\text{SiO}_2 \cdot 3.59\text{CH}_4 \cdot 3.10\text{N}_2 \cdot 1.31\text{CO}_2$ from Racalmuto, Sicily, along with a heat-treated (molecule-free) sample of composition SiO_2 , were studied between 5 and 300 K using heat-pulse microcalorimetry. The molecule-free sample was obtained by heating natural Racalmuto crystals at 1173 K for 24 h. The standard third-law entropy of the molecule-free sample is $S^\circ = 2216.3 \pm 6.6 \text{ J}/(\text{mol} \cdot \text{K})$ for 46SiO_2 and the natural Mt. Hamilton and Racalmuto samples give $S^\circ = 2805.7 \pm 8.4 \text{ J}/(\text{mol} \cdot \text{K})$ and $S^\circ = 2956.8 \pm 8.9 \text{ J}/(\text{mol} \cdot \text{K})$, respectively. The entropy and Gibbs free energy for molecule-free melanophlogite relative to quartz at 298 K are $\Delta S_{\text{trans}} = 6.7 \text{ J}/(\text{mol} \cdot \text{K})$ and $\Delta G_{\text{trans}} = 7.5 \text{ kJ}/\text{mol}$, respectively and, thus, it does not have a thermodynamic field of stability in the SiO_2 system. The difference in C_p values between molecule-containing and molecule-free melanophlogite is characterized by an increase in C_p from 0 to $\sim 70 \text{ K}$, and it then reaches a roughly constant value at $70 \text{ K} < T < 250 \text{ K}$. The ΔS^{rxn} at 298 K for $46\text{SiO}_2(\text{melan.}) + x\text{CH}_4(\text{gas}) + y\text{CO}_2(\text{gas}) + z\text{N}_2(\text{gas}) = 46\text{SiO}_2 \cdot (x\text{CH}_4)^{12} \cdot (y\text{CO}_2, z\text{N}_2)^{14}$ is estimated to be about -642 and $-802 \text{ J}/(\text{mol} \cdot \text{K})$ for the Mt. Hamilton and Racalmuto samples, respectively. The thermodynamic data, as well as published results on the occurrence of natural molecule-containing samples suggest that melanophlogite crystallizes metastably. The occurrence of melanophlogite and the lack of other porosils in nature are probably due to the essential role of molecular structure-directing agents. For melanophlogite they can be CO_2 , N_2 , and CH_4 , whereas the crystallization of other porosils requires more chemically and structurally complex molecules that are not naturally abundant.

Keywords: Melanophlogite, heat capacity, entropy, clathrasils, microporous minerals, clathrate