

## **Dmitryivanovite: A new high-pressure calcium aluminum oxide from the Northwest Africa 470 CH3 chondrite characterized using electron backscatter diffraction analysis**

**TAKASHI MIKOUCHI,<sup>1,\*</sup> MICHAEL ZOLENSKY,<sup>2</sup> MARINA IVANOVA,<sup>3</sup> OSAMU TACHIKAWA,<sup>1</sup>  
MUTSUMI KOMATSU,<sup>1</sup> LOAN LE,<sup>4</sup> AND MATTHIEU GOUNELLE<sup>5</sup>**

<sup>1</sup>Department of Earth and Planetary Science, Graduate School of Science, University of Tokyo, Hongo, Bunkyo-Ku, Tokyo 113-0033, Japan

<sup>2</sup>Code KT, NASA Johnson Space Center, Houston, Texas 77058, U.S.A.

<sup>3</sup>Vernadsky Institute of Geochemistry, Kosygin Street 19, Moscow 119991, Russia

<sup>4</sup>Jacobs Sverdrup Co., Houston, Texas 77058, U.S.A.

<sup>5</sup>Laboratoire d'Étude, de la Matière Extraterrestre, Muséum National d'Histoire Naturelle, 57 rue Cuvier, 75005 Paris cedex, France

### **ABSTRACT**

Dmitryivanovite ( $\text{CaAl}_2\text{O}_4$ ) is a newly described, calcium aluminum oxide from the Northwest Africa 470 (NWA470) CH3 chondrite (Ivanova et al. 2002). NWA470 contains abundant small Ca,Al-rich inclusions (CAIs), and dmitryivanovite, whose composition is close to stoichiometric  $\text{CaAl}_2\text{O}_4$  [ $\text{Ca}_{1.000}(\text{Al}_{1.993}\text{Si}_{0.003}\text{Ti}_{0.002})_{1.998}\text{O}_4$ ], was found in one of these CAIs. It occurs as  $\sim 10\ \mu\text{m}$  subhedral grains intergrown with grossite ( $\text{CaAl}_4\text{O}_7$ ), perovskite, and melilite. Electron backscatter diffraction (EBSD) analysis revealed that dmitryivanovite is a high-pressure polymorph of  $\text{CaAl}_2\text{O}_4$  ( $a = 7.95$ ,  $b = 8.62$ ,  $c = 10.25\ \text{Å}$ ,  $\beta = 93.1^\circ$ , space group  $P2_1/c$ , and  $Z = 12$ ). Dmitryivanovite is the third phase to be described from nature in the binary system of  $\text{CaO}-\text{Al}_2\text{O}_3$ , the other two being hibonite ( $\text{CaAl}_{12}\text{O}_{19}$ ) and grossite ( $\text{CaAl}_4\text{O}_7$ )—all are found in CAIs. The presence of  $\text{CaAl}_2\text{O}_4$  in NWA470 suggests a local elevated dust/gas ratio in the solar nebula. The phase diagram of  $\text{CaAl}_2\text{O}_4$  shows that  $\sim 2\ \text{GPa}$  is required to stabilize the high-pressure  $\text{CaAl}_2\text{O}_4$  polymorph at  $1327\ ^\circ\text{C}$ , above which  $\text{CaAl}_2\text{O}_4$  condenses from the solar nebula. Because it is unlikely that the solar nebula ever had such a high total gas pressure, it appears more probable that condensation of the low-pressure polymorph occurred in the solar nebula with an enhanced dust-to-gas ratio and that subsequently the high-pressure polymorph was produced by shock metamorphism, most likely after the  $\text{CaAl}_2\text{O}_4$ -bearing CAI was incorporated into the NWA470 parent asteroid.

**Keywords:** Dmitryivanovite, CAI, electron backscatter diffraction, new minerals, CH chondrite