

LETTER

The structure of a super-aluminous version of the dense hydrous-magnesium silicate phase D

TIZIANA BOFFA BALLARAN,* DANIEL J. FROST, NOBUYOSHI MIYAJIMA, AND FLORIAN HEIDELBACH

Bayerisches Geoinstitut, Universität Bayreuth, D-95440 Bayreuth, Germany

ABSTRACT

The dense hydrous-magnesium silicate phase D, which has the ideal formula $\text{MgSi}_2\text{H}_2\text{O}_6$, may be an important link in a chain of hydrous phases that carry H_2O in the ultramafic portions of subducting lithosphere, into the Earth's lower mantle. We have synthesized a new Al-rich form of phase D, containing up to 50 wt% Al_2O_3 , using a multi-anvil device at $\sim 1300^\circ\text{C}$ and 25 GPa. The phase, with the formula $\text{Mg}_{0.2}\text{Fe}_{0.15}\text{Al}_{1.8}\text{H}_{1.8}\text{SiO}_6$, was initially produced in a bulk composition designed to synthesize Al- and Fe-rich magnesium silicate perovskite with a composition similar to that produced in experiments on mid-ocean ridge basalt bulk compositions at lower mantle conditions. Further experiments using a starting mixture based on the composition of this Al-rich phase resulted in the synthesis of 60–70 μm long single crystals at similar conditions. The recovered crystals were slightly richer in H_2O ($\text{Mg}_{0.2}\text{Fe}_{0.12}\text{Al}_{1.5}\text{Si}_{0.92}\text{H}_{3.1}\text{O}_6$) and their unit-cell parameters were similar to those of $\text{MgSi}_2\text{H}_2\text{O}_6$ phase D. A refinement of the crystal structure was carried out in the $P\bar{3}1m$ space group and revealed a more disordered cation distribution than magnesium silicate phase D. All cation-oxygen distances are similar, suggesting a high degree of Si/Al disorder. Although the stability field of this new variant of phase D is yet to be determined, this phase may be an important host for H_2O within portions of subducted oceanic crust in the lower mantle.

Keywords: Lower mantle, high pressure, subduction, single crystal, DHMS, deep water cycle