

## Coexisting Actinolite and Hornblende from West-Central New Hampshire

JOHN B. BRADY

*Department of Geological Sciences, Harvard University  
Cambridge, Massachusetts 02138*

### Abstract

The coexistence of actinolite and hornblende and of cummingtonite and hornblende has been observed in garnet-zone and staurolite-kyanite-zone amphibolites of the Ammonoosuc Volcanics near Hanover, New Hampshire. The amphibole pairs occur both as separate grains and as intergrowths with boundaries that are optically and chemically sharp, but geometrically irregular; no exsolution lamellae were observed. Electron microprobe analyses of the amphibole pairs show that: (1) hornblende is richer in Al, Na, K, Ti, and Fe<sup>3+</sup>, and has a lower Mg/(Mg + Fe) ratio than the coexisting actinolite; (2) hornblende is enriched in Al, Na, and Ca, but has a similar Mg/(Mg + Fe) ratio to that of the coexisting cummingtonite. Recalculations show that the A site is nearly empty in actinolites and typically one-third to one-half full in coexisting hornblendes. Single-crystal precession photos of actinolite-hornblende crystals show that the lattices of the two phases share a common *b*-axis, but are rotated relative to one another by two to three degrees.

### Introduction

The existence of a miscibility gap in the calcic amphiboles has been substantiated only recently (Hallimond, 1943; Shido and Miyashiro, 1959; Klein, 1969), probably because of the optical similarity between coexisting actinolite and hornblende. In addition, the two amphiboles are commonly found together in a patchy intergrowth that may be easily overlooked, rather than as single grains showing exsolution lamellae. Klein (1969) and Cooper and Lovering (1970) have presented analyses for nine actinolite-hornblende pairs from a variety of localities. Although their analyses reveal several important features of the miscibility gap, there are too many variables involved to explain the variety of compositions observed. This report presents new analyses of actinolite-hornblende pairs from several rocks of a single metamorphic grade. It is a first step toward a more detailed understanding of the calcic amphibole miscibility gap.

### Geologic Setting of Amphibole Pairs

All the rocks studied were collected from the Middle Ordovician Ammonoosuc Volcanics in west-central New Hampshire. Specifically, all the amphibole pairs are from cliff-like outcrops on the west flank of the Mascoma gneiss dome, primarily in the

Mascoma Quadrangle (location map in Brady, 1970). Details of the local geology may be found in Chapman (1939), Hadley (1942), and Naylor (1968). The sample localities closely straddle a staurolite isograd, samples coming mainly from the staurolite-kyanite zone (Thompson *et al.*, 1968).

The Ammonoosuc Volcanics consist primarily of fine-grained amphibolites with subordinate felsic layers. The formation is believed to be volcanic in origin. Billings (1937, p. 476) has described remnant volcanic structures in the less-metamorphosed Ammonoosuc of the Littleton-Moosilauke area, and Chapman (1939, p. 135) found structures resembling those of volcanic tuffs and breccias in the area studied here. The amphibolites contain hornblende and plagioclase as the principal minerals, though epidote is locally abundant. Estimated modes of the polished sections used in electron probe microanalysis are given in Table 1.

Coexisting amphiboles were distinguished primarily on the basis of color and pleochroism. The hornblende is pleochroic deep green to blue-green. Coexisting actinolite is nearly colorless, slightly pleochroic from pale blue-green to green. When more iron-rich, the amphiboles are more highly colored. A difference in birefringence commonly accompanies the color difference, the actinolite having the higher order interference colors. Where cummingtonite coexists

TABLE 1. Estimated Modes of Samples Containing Amphibole Pairs\*

	C-2	G	C-6	17
hornblende	32	28	77	18
actinolite	7	15	6	--
cummingtonite	--	--	--	13
epidote	40	47	--	--
chlorite	--	--	10	7
plagioclase	20 (An <sub>50</sub> )	3 (An <sub>30</sub> )	2 (An <sub>15</sub> )	16 (An <sub>25</sub> )
quartz	--	4	3	45
sphene	1	3	--	--
rutile	--	tr.**	2	--
zircon	--	--	--	tr.
biotite	--	--	--	tr.
ilmenite	--	--	--	1

\* numbers are given in volume percent

\*\* tr. = trace, much less than one percent

with hornblende, its colorless crystals and optically positive character permit easy identification.

Contacts between the coexisting amphiboles are optically sharp but may have an irregular shape (Fig. 1A). In some occurrences the two amphiboles appear to form a single crystal that is a "patchwork" of the two phases (see Klein, 1969, Figure 1). Actinolite-hornblende pairs commonly exhibit this




FIG. 1. A. Photomicrograph of sample #C2 in plain-polarized light showing actinolite-hornblende pairs. The grass-green hornblende is the darker mineral. The very pale green actinolite looks colorless in the photo. Note the optically sharp contacts between the two amphiboles which occur both as separate grains and as "patchy" intergrowths. The high relief mineral is epidote; the low relief mineral is plagioclase. The field of view is  $0.8 \times 0.8$  mm.

relationship. Cummingtonite-hornblende intergrowths are more regular, the two phases sharing ends or sides of the same crystal (Figure 1B). Extinction of both phases of these two-amphibole crystals appears simultaneous. Whether or not these textures are consistent with the postulate of chemical equilibrium between the two amphiboles is an important question which has been addressed by others (Klein, 1969; Stout, 1971). I believe that the presence of a sharp optical and chemical discontinuity between two homogeneous amphiboles is certainly consistent with equilibrium and strong evidence in favor of a miscibility gap. No exsolution lamellae were observed in any of the amphiboles.

### Chemical Analyses

Amphibole pairs were analyzed in polished carbon-coated thin sections on an Applied Research Laboratories (EMX) microprobe. Nine element analyses were obtained using the methods of Bence and Albee (1968) and Albee and Ray (1970) for an accelerating potential of 15 kV. Seven element analyses were obtained using the standard curve method as described by Klein (1968) for an accelerating potential of 20 kV. The analyses listed in Table 2 have been




FIG. 1. B. Photomicrograph of sample #17 in plain-polarized light showing cummingtonite-hornblende pairs in a quartz matrix. The hornblende is pale green; the cummingtonite is colorless. Note particularly the crystals which have hornblende at both ends with cummingtonite between. The field of view is  $0.6 \times 0.6$  mm.

TABLE 2. Chemical Analyses of Actinolite-Hornblende and Cummingtonite-Hornblende Pairs

Hornblende	Sample #C2					Sample #C					Sample #C6			Sample #17	
	D	E	H	I	Z	H	H	H	J1	C	D	F	Z	4	
SiO <sub>2</sub>	45.40	46.28	45.21	45.4	44.7	43.06	46.00	45.06	45.00	45.21	47.09	44.45	46.8	46.9	
Al <sub>2</sub> O <sub>3</sub>	9.72	9.52	10.36	10.8	11.0	13.66	11.28	12.01	12.30	13.11	11.67	13.94	11.1	10.8	
TiO <sub>2</sub>	0.45	0.47	0.44	n.d.*	n.d.*	0.56	0.47	0.48	0.55	0.58	0.39	0.57	n.d.	n.d.	
FeO	18.03	17.53	18.45	17.0	18.6	15.47	13.71	15.16	14.39	10.31	10.30	10.60	13.5	13.4	
MnO	0.43	0.34	0.46	0.3	0.4	0.54	0.58	0.49	0.52	0.20	0.23	0.22	0.3	0.3	
MgO	9.96	10.36	9.58	9.7	9.1	10.45	12.34	11.33	11.73	13.93	14.36	13.20	13.0	12.9	
CaO	12.09	12.10	11.97	12.3	12.3	11.89	12.13	12.22	12.11	10.83	11.25	10.87	11.1	11.0	
Na <sub>2</sub> O	1.29	1.29	1.29	1.1	1.2	2.03	1.66	1.73	1.76	1.90	1.84	2.11	1.7	1.6	
K <sub>2</sub> O	0.32	0.29	0.38	n.d.	n.d.	0.24	0.17	0.17	0.15	0.14	0.12	0.21	n.d.	n.d.	
Total	97.63	98.12	98.14	96.6	97.3	97.88	98.34	98.65	98.31	96.21	97.25	96.17	97.5	96.9	
Si **	6.765 (15)***	6.860 (20)	6.715 (25)	6.785 (5)	6.675 (5)	6.345 (25)	6.670 (30)	6.560 (20)	6.535 (25)	6.540 (60)	6.745 (65)	6.465 (65)	6.765 (65)	6.810 (60)	
Al (IV)	1.235 (15)	1.160 (20)	1.285 (25)	1.215 (5)	1.325 (5)	1.655 (25)	1.330 (30)	1.440 (20)	1.465 (25)	1.480 (60)	1.255 (65)	1.535 (65)	1.235 (65)	1.190 (60)	
Al (VI)	0.470 (20)	0.495 (25)	0.530 (30)	0.685 (5)	0.610 (5)	0.715 (35)	0.600 (40)	0.620 (30)	0.640 (30)	0.780 (70)	0.720 (50)	0.855 (65)	0.655 (85)	0.660 (80)	
Ti	0.05 (0)	0.05 (0)	0.05 (0)	n.d.	n.d.	0.06 (0)	0.05 (0)	0.05 (0)	0.06 (0)	0.06 (0)	0.04 (0)	0.06 (0)	n.d.	n.d.	
Fe <sup>+3</sup>	0.310 (130)	0.245 (135)	0.315 (145)	0.240 (50)	0.395 (45)	0.310 (200)	0.250 (190)	0.290 (150)	0.300 (190)	0.405 (405)	0.300 (300)	0.340 (340)	0.425 (425)	0.415 (415)	
Fe <sup>+2</sup>	1.935 (135)	1.925 (145)	1.970 (160)	1.890 (50)	1.930 (50)	1.595 (205)	1.415 (195)	1.560 (160)	1.445 (195)	0.845 (415)	0.935 (305)	0.950 (350)	1.235 (65)	1.210 (60)	
Mn	0.05 (0)	0.04 (0)	0.06 (0)	0.04 (0)	0.05 (0)	0.07 (0)	0.07 (0)	0.06 (0)	0.06 (0)	0.06 (0)	0.02 (0)	0.03 (0)	0.04 (0)	0.04 (0)	
Mg	2.215 (5)	2.285 (5)	2.120 (10)	2.16 (0)	2.025 (5)	2.300 (10)	2.670 (10)	2.460 (10)	2.540 (10)	3.005 (25)	3.070 (20)	2.860 (20)	2.800 (30)	2.795 (25)	
Ca	1.925 (5)	1.915 (5)	1.905 (5)	1.97 (0)	1.97 (0)	1.880 (10)	1.885 (5)	1.905 (5)	1.885 (5)	1.675 (15)	1.730 (10)	1.695 (15)	1.715 (15)	1.715 (15)	
Na (M4)	0.045 (30)	0.045 (40)	0.050 (50)	0.015 (15)	0.020 (20)	0.070 (60)	0.060 (60)	0.055 (50)	0.070 (50)	0.210 (130)	0.180 (100)	0.210 (110)	0.155 (145)	0.145 (130)	
Na (A)	0.325 (30)	0.305 (40)	0.320 (50)	0.305 (15)	0.305 (15)	0.510 (60)	0.405 (65)	0.435 (50)	0.425 (50)	0.325 (135)	0.385 (115)	0.385 (150)	0.290 (150)	0.290 (130)	
K	0.06 (0)	0.05 (0)	0.07 (0)	n.d.	n.d.	0.045 (5)	0.03 (0)	0.03 (0)	0.03 (0)	0.03 (0)	0.02 (0)	0.04 (0)	n.d.	n.d.	
Sum Mg	13.030 (30)	13.040 (40)	13.045 (45)	13.015 (15)	13.010 (10)	13.050 (50)	13.055 (55)	13.060 (60)	13.045 (65)	13.115 (115)	13.090 (90)	13.095 (95)	13.130 (130)	13.125 (115)	
Sum K	15.385 (45)	15.355 (45)	15.390 (50)	15.305 (15)	15.330 (20)	15.555 (65)	15.435 (65)	15.465 (45)	15.455 (55)	15.355 (135)	15.350 (100)	15.425 (115)	15.320 (150)	15.290 (130)	
Actinolites															
SiO <sub>2</sub>	54.32	54.05	54.64	54.6	54.0	53.03	52.91	53.56	53.67	54.52	53.73	54.01	53.8	54.1	
Al <sub>2</sub> O <sub>3</sub>	1.67	2.14	1.45	1.5	2.2	4.28	4.32	3.77	4.21	2.16	3.46	2.93	0.6	0.2	
TiO <sub>2</sub>	0.05	0.05	0.03	n.d.	n.d.	0.15	0.11	0.09	0.13	0.06	0.12	0.05	n.d.	n.d.	
FeO	13.44	13.53	13.80	13.2	13.4	11.88	10.87	10.87	10.44	8.11	9.08	8.37	20.3	21.0	
MnO	0.38	0.37	0.37	0.3	0.3	0.45	0.47	0.46	0.45	0.26	0.25	0.28	1.0	1.1	
MgO	15.30	15.08	15.05	15.3	15.1	15.19	16.19	16.48	16.67	18.97	17.95	18.53	18.4	18.0	
CaO	12.73	12.74	12.78	12.8	12.7	12.37	12.61	12.68	12.69	11.98	12.21	12.04	0.9	0.8	
Na <sub>2</sub> O	0.18	0.19	0.13	0.1	0.2	0.70	0.57	0.45	0.53	0.32	0.50	0.44	0.1	0.0	
K <sub>2</sub> O	0.06	0.04	0.06	n.d.	n.d.	0.07	0.04	0.03	0.03	0.04	0.06	0.09	n.d.	n.d.	
Total	98.13	98.19	98.31	97.8	97.9	98.12	98.09	98.39	98.82	96.42	97.36	96.74	95.1	95.2	
Si	7.800 (10)	7.760 (10)	7.845 (5)	7.855 (5)	7.765 (15)	7.575 (15)	7.520 (20)	7.580 (20)	7.550 (20)	7.750 (20)	7.600 (30)	7.660 (30)	7.940 (40)	8.005 (35)	
Al (IV)	0.200 (10)	0.240 (10)	0.155 (5)	0.145 (5)	0.235 (15)	0.425 (15)	0.480 (20)	0.420 (20)	0.450 (20)	0.250 (20)	0.400 (30)	0.340 (30)	0.060 (40)	0.035 (5)	
Al (VI)	0.080 (10)	0.120 (10)	0.095 (5)	0.105 (5)	0.135 (15)	0.295 (15)	0.245 (25)	0.210 (20)	0.250 (20)	0.110 (20)	0.115 (35)	0.185 (30)	0.040 (40)	0.035 (5)	
Ti	0.01 (0)	0.01 (0)	0.00 (0)	n.d.	n.d.	0.02 (0)	0.01 (0)	0.01 (0)	0.01 (0)	0.01 (0)	0.01 (0)	0.01 (0)	n.d.	n.d.	
Fe <sup>+3</sup>	0.095 (65)	0.095 (65)	0.050 (50)	0.040 (40)	0.090 (70)	0.065 (65)	0.130 (130)	0.145 (125)	0.130 (130)	0.130 (130)	0.195 (195)	0.170 (170)	0.255 (255)	0.215 (215)	
Fe <sup>+2</sup>	1.520 (70)	1.530 (70)	1.605 (55)	1.545 (45)	1.520 (70)	1.350 (70)	1.165 (135)	1.140 (130)	1.100 (130)	0.840 (130)	0.880 (200)	0.825 (175)	2.250 (270)	2.380 (230)	
Mn	0.05 (0)	0.045 (5)	0.045 (5)	0.04 (0)	0.04 (0)	0.05 (0)	0.06 (0)	0.05 (5)	0.05 (0)	0.03 (0)	0.03 (0)	0.03 (0)	0.125 (5)	0.14 (0)	
Mg	3.275 (5)	3.225 (5)	3.22 (0)	3.28 (0)	3.235 (5)	3.235 (5)	3.430 (10)	3.480 (10)	3.495 (5)	4.020 (10)	3.785 (15)	3.915 (15)	4.050 (20)	3.980 (10)	
Ca	1.96 (0)	1.96 (0)	1.965 (5)	1.97 (0)	1.955 (0)	1.895 (5)	1.925 (5)	1.925 (5)	1.915 (5)	1.825 (5)	1.850 (10)	1.830 (10)	0.16 (0)	0.13 (0)	
Na (M4)	0.010 (10)	0.015 (15)	0.020 (20)	0.020 (10)	0.025 (25)	0.090 (20)	0.035 (35)	0.035 (35)	0.050 (40)	0.035 (35)	0.075 (65)	0.070 (60)	0.030 (30)	0.00 (0)	
Na (A)	0.040 (10)	0.035 (15)	0.020 (20)	0.010 (10)	0.035 (25)	0.100 (20)	0.125 (35)	0.085 (40)	0.090 (40)	0.055 (45)	0.065 (65)	0.050 (60)	n.d.	n.d.	
K	0.01 (0)	0.01 (0)	0.01 (0)	n.d.	n.d.	0.01 (0)	0.01 (0)	0.01 (0)	0.01 (0)	0.01 (0)	0.01 (0)	0.02 (0)	n.d.	n.d.	
Sum Mg	13.030 (30)	13.025 (25)	13.015 (15)	13.010 (10)	13.020 (20)	13.015 (15)	13.040 (40)	13.040 (40)	13.035 (35)	13.140 (40)	13.075 (55)	13.100 (50)	14.720 (80)	14.755 (65)	
Sum K	15.050 (20)	15.045 (25)	15.030 (20)	15.010 (10)	15.035 (25)	15.110 (20)	15.135 (45)	15.095 (45)	15.100 (40)	15.065 (45)	15.075 (65)	15.070 (60)	14.890 (80)	14.885 (65)	

\* n.d. means not determined. The seven element analyses were completed using standard curves (Klein, 1968). The nine element analyses were completed using empirical correction factors (Bence and Albee, 1968).

\*\* See text for description of recalculation scheme. Also see Stout (1972). Sums given are cumulative down the column of formula unit numbers.

\*\*\* Parenthesized figures represent one half the difference between the greatest lower bound and least upper bound recalculations (see text) in terms of least units cited for the value to the immediate left, thus 6.765 (15) means that the value 6.765 is in the middle of a bracket 0.030 units wide. The "true" value could be anywhere within the bracket.

selected as representative of the 35 amphibole pairs actually analyzed.

The chemical analyses have been recalculated utilizing chemical and stoichiometric constraints to give upper and lower bounds for the value of each cation in the formula unit. This type of recalculation is necessitated by the indeterminacy of the valence state of iron in electron microprobe analyses. It is common practice simply to consider all the iron to be ferrous. However, it is perhaps more useful to use the knowledge available concerning a mineral's structure and stoichiometry to determine maximum and minimum values for ferric iron, between which the "real" value lies. The range of values so determined decreases in size as the iron-content of the mineral increases. A more complete discussion of this type of recalculation procedure is given in Stout (1972).

All recalculations used the following assumptions:

- (1) there are 23 oxygens exclusive of H<sub>2</sub>O per half unit cell;
- (2) the analyses are complete except for H<sub>2</sub>O;
- (3) there are no vacancies except in the A site. One additional assumption is sufficient to specify the ferric-ferrous ratio. For the calcic amphiboles the most useful numbers are provided by assumptions concerning the occupancy of the M(4) site. Usually, the tightest brackets are obtained by assuming that there is no sodium or potassium in the M(4) site (minimum Fe<sup>3+</sup>) or that there is only calcium and sodium in the M(4) site (maximum Fe<sup>3+</sup>). In some cases, the assumption of all ferrous iron (minimum Fe<sup>3+</sup>) or the assumption that there is only potassium in the A site (maximum Fe<sup>3+</sup>) will provide more restrictive bounds. Results of computer calculations are presented in Table 2 along with the chemical analyses. The numbers given are the mean of the greatest lower bound and least upper bound recalculated for each

analysis. The numbers in parentheses are half the difference of the two bounds in terms of the least units cited for the value to the immediate left.

For the actinolite-hornblende pairs the analyses show the following element fractionations: (1) Actinolite is richer in Mg and Si and very poor in Al, Na, and K. (2) Hornblende is richer in Al, Na, K, Ti, and  $\text{Fe}^{3+}$  (calc) and has a lower  $\text{Mg}/(\text{Mg} + \text{Fe})$  ratio. The cation sums below the recalculations show that in actinolites the *A* site is nearly, but not quite, empty. The coexisting hornblendes typically yield values for the *A* site one-third to one-half full.

The actinolite varies from an ideal two-component composition (tremolite-ferroactinolite) principally utilizing the tschermakite  $[\text{Al}^{\text{VI}}\text{Al}^{\text{IV}} \rightleftharpoons (\text{Mg}, \text{Fe}^{2+})^{\text{VI}}\text{Si}^{\text{IV}}]$  and edenite  $[\text{Na}^{\text{A}}\text{Al}^{\text{IV}} \rightleftharpoons \square^{\text{A}}\text{Si}^{\text{IV}}]$  substitutions (Fig. 2).<sup>1</sup> The hornblende, however, has a significant

<sup>1</sup> The superscripts refer to coordination number. *A* refers to the *A* site, which has 10-12-fold coordination. *M*(4) refers to the *M*(4) site, which has 6-8-fold coordination.  $\square$  means a vacancy.

$[(\text{Fe}^{3+})^{\text{VI}} \rightleftharpoons \text{Al}^{\text{VI}}]$  substitution as well, tending toward a magnesio-hastingsite composition. A few of the hornblendes also have a noticeable glaucophane component  $[\text{Na}^{\text{M}(4)}\text{Al}^{\text{VI}} \rightleftharpoons \text{Ca}^{\text{M}(4)}(\text{Mg}, \text{Fe}^{2+})^{\text{VI}}]$ . Interestingly, Figure 3 shows that alkali content is higher for amphiboles with a higher  $\text{Mg}/(\text{Mg} + \text{Fe})$  ratio, particularly for hornblendes. The tschermakite component also increases with the  $\text{Mg}/(\text{Mg} + \text{Fe})$  ratio. However, these effects may be an accident of bulk composition. In general, the compositions observed are consistent with the trends discussed by Robinson, Ross, and Jaffe (1971a) for the calcic amphiboles and related anthophyllite-gedrite series.

Analyses of coexisting cummingtonite and hornblende show element fractionations consistent with the results of more detailed work by others (Klein, 1968; Robinson and Jaffe, 1969, with wet analyses of two pairs). (1) Cummingtonite is richer in Mg, Mn, Fe, and Si. (2) Hornblende is richer in Ca, Al, and Na and has a slightly higher  $\text{Mg}/(\text{Mg} + \text{Fe})$




FIG. 2. The actinolite-hornblende pairs are shown here on a compositional plane which approximates the end-member system tremolite-edenite-tschermakite. The ordinate is the calculated *A* site occupancy per half unit cell (total number of cations minus 15.00). The abscissa is the calculated quantity of tetrahedral aluminum (8.00 minus the number of silicon atoms per half unit cell) minus the calculated *A* site occupancy. These axes were chosen to make the end-member system orthogonal. The dashed tie lines are for pairs for which potassium was not determined.

ratio. Totals for the cummingtonite analyses are low, perhaps because of the standard curves used. The cummingtonite-hornblende data is presented primarily because these pairs were found within meters of the actinolite-hornblende pairs (Fig. 4).

### X-ray Crystallography

Selected actinolite-hornblende crystals from several samples were examined using single-crystal precession and rotation X-ray techniques to determine: (1) the possible presence of amphiboles with a primitive monoclinic lattice, (2) the unit-cell parameters of the coexisting phases, and (3) the relative orientations of the amphibole lattices in "two-phase" crystals. Crystals were picked from samples within millimeters of the area of the analyzed amphibole pairs. The "two-phase" crystals chosen, although part green, part colorless, appeared otherwise structurally continuous.

Of the crystals photographed, none showed violations of  $C2/m$  symmetry. There did appear to be a consistent relationship in the orientations of the




FIG. 3. Compositions of the actinolite-hornblende pairs are shown here in a comparison of alkali content and magnesium to magnesium plus total iron ratio. The hornblendes have a significant alkali content lacking in coexisting actinolites. The alkali content of the hornblendes is greater for hornblendes with a higher magnesium to magnesium plus total iron ratio. The dashed tie lines are for pairs for which potassium was not determined.

TABLE 3. Unit Cell Parameters for "Two-Phase" Actinolite-Hornblende Crystals\*

	Crystal C-2b		Crystal C-6b	
	Hornblende	Actinolite	Hornblende	Actinolite
$a$ (Å)	9.84	9.86	9.83	9.81
$b$ (Å)	18.12	n.d.**	18.07	n.d.
$c$ (Å)	5.30	5.28	5.30	5.30
$\beta$	$104^{\circ}52'$	$104^{\circ}48'$	$104^{\circ}55'$	$104^{\circ}54'$
Space group	$C2/m$	$C2/m$	$C2/m$	$C2/m$

\* Results of measurements of precession photos. Cell edges are precise to  $\pm 0.5$  percent, angular values to  $\pm 3'$ .

\*\* n.d. = not determined

lattices of the actinolite-hornblende intergrowths. Three different actinolite-hornblende crystals showed lattices orientated within 2 to 3 degrees of rotation (+ or -) about a common  $b$  axis. The  $b$  axes were very nearly, though not exactly, coincident. The unit cells of the two phases are quite similar (Table 3).

The orientation of exsolution lamellae in hornblende-cummingtonite crystals has been discussed by Robinson *et al* (1971b). These authors conclude that the two amphibole phases are interfaced along planes of "best fit" of the two lattices. The irregular intergrowths of actinolite and hornblende observed in this study, while possibly formed by a different process, clearly are not interfaced on a plane of best fit. The intergrowths do show a close relative lattice orientation, however. It follows that one of the amphibole phases (or a homogeneous parent) pre-dates the other and that the orientation of the second amphibole lattice was controlled by that of the first. The mechanism of this structural control may involve a nucleation or a replacement process.

The textures presented by patchy, two-amphibole crystals are most intriguing. Why are the contacts between the two phases commonly so irregular? Such boundaries seem to be far from representing a minimization of the potential energy of the interface. It may be that the assemblages themselves are a long way from being equilibrium assemblages. On the other hand, it may be that the potential energy per unit area of interface is very low, due to the similarity of the unit cells of the phases involved. If so, then there is little to be gained, thermodynamically, by a more regular geometry. Evidence in favor of the latter view may be found in an examination of the list of amphibole pairs that exhibit the irregular intergrowth texture. This list includes actinolite-hornblende, glaucophane-hornblende, actinolite-glaucophane (Klein, 1969), and anthophyllite-gedrite




FIG. 4. Compositions of both actinolite-hornblende pairs and cummingtonite-hornblende pairs are shown on a portion of the triangle  $\text{CaO}-(\text{Al}_2\text{O}_3-\text{Na}_2\text{O})$ ,  $\text{MgO}$ ,  $\text{FeO} + \text{MnO}$ , which is the plagioclase projection of Robinson and Jaffe (1969). This diagram conveniently shows relationships between the compositions of the cummingtonite-hornblende pairs (found within meters of actinolite-hornblende sample #G) and the compositions of the actinolite-hornblende pairs.

(Stout, 1971)<sup>2</sup>. Noticeably absent from the list are pairs consisting of cummingtonite and a calcic or sodic amphibole, calcic amphibole-orthoamphibole pairs, or sodic amphibole-orthoamphibole pairs. These are the combinations which have the greatest contrast in structure and, therefore, possibly the greatest potential energy reduction due to a regulated interface geometry.

#### Acknowledgments

This report summarizes the content of work submitted as a Harvard College undergraduate thesis in 1970. Stimulus for and guidance throughout were provided by my advisors Professors James B. Thompson, Jr., Charles W. Burnham, and Cornelis Klein, Jr. I am also indebted to James H. Stout, Joel E. Arem, and Richard Beger who neglected their own educations in the interests of mine. Peter Robinson made several suggestions that improved the final manuscript. The chemical analytical work was supported by NSF Grant GA-11435 (Thompson and Klein). The final preparation was completed while the author held an NSF Graduate Fellowship.

#### References

- ALBEE, A. L., AND L. RAY (1970) Correction factors for electron probe microanalysis of silicates, oxides, carbonates, phosphates, and sulphates. *Anal. Chem.* **42**, 1408-1414.
- BENCE, A. E., AND A. L. ALBEE (1968) Empirical correction factors for the electron microanalysis of silicates and oxides. *J. Geol.* **76**, 382-403.
- BILLINGS, M. P. (1937) Regional metamorphism of the Littleton-Moosilauke area, New Hampshire. *Bull. Geol. Soc. Am.* **48**, 463-566.
- BRADY, JOHN B. (1970) *Amphibole Assemblages from the Ammonoosuc Volcanics of the Mascoma Dome*. A.B. Thesis, Harvard College, Cambridge, Massachusetts.
- CHAPMAN, C. A. (1939) Geology of the Mascoma quadrangle, New Hampshire. *Bull. Geol. Soc. Am.* **50**, 127-180.
- COOPER, A. F., AND J. F. LOVERING (1970) Greenschist amphiboles from Haast River, New Zealand. *Contrib. Mineral. Petrol.* **27**, 11-24.
- HADLEY, J. B. (1942) Stratigraphy, structure, and petrology of the Mt. Cube area, New Hampshire. *Bull. Geol. Soc. Am.* **53**, 113-176.
- HALLIMOND, A. F. (1943) On the graphical representation of the calciferous amphiboles. *Am. Mineral.* **28**, 65-89.
- KLEIN, CORNELIS, JR. (1968) Coexisting amphiboles. *J. Petrol.* **9**, 281-330.
- (1969) Two-amphibole assemblages in the system actinolite-hornblende-glaucophane. *Am. Mineral.* **54**, 212-237.
- NAYLOR, R. S. (1968) Origin and regional relationships of the core-rocks of the Oliverian domes. In E-an Zen, W. S. White, J. B. Hadley, and J. B. Thompson, Jr., Eds., *Studies of Appalachian Geology, Northern and Maritime*. John Wiley and Sons, New York, p. 231-240.

<sup>2</sup> Some of these pairs, notably anthophyllite-gedrite, also occur as regular intergrowths.

- ROBINSON, PETER, AND H. W. JAFFE (1969) Chemographic exploration of amphibole assemblages from Central Massachusetts and southwestern New Hampshire. *Mineral. Soc. Am. Spec. Pap.* **2**, 251-274.
- , MALCOLM ROSS, AND H. W. JAFFE (1971a) Composition of the anthophyllite-gedrite series, comparisons of gedrite and hornblende, and the anthophyllite-gedrite solvus. *Am. Mineral.* **56**, 1005-1041.
- , H. W. JAFFE, MALCOLM ROSS, AND CORNELIS KLEIN, JR. (1971b) Orientation of exsolution lamellae in clinopyroxenes and clinoamphiboles: Consideration of optimal phase boundaries. *Am. Mineral.* **56**, 909-939.
- SHIDO, FUMIKO, AND AKIHO MIYASHIRO (1959) Hornblendes of basic metamorphic rocks. *J. Fac. Sci. Univ. Tokyo*, **12**, 85-102.
- STOUT, JAMES H. (1971) Four coexisting amphiboles from Telemark, Norway. *Am. Mineral.* **56**, 212-224.
- (1972) Phase petrology and mineral chemistry of coexisting amphiboles from Telemark, Norway, *J. Petrol.* **13**, 99-145.
- THOMPSON, J. B., JR., PETER ROBINSON, T. N. CLIFFORD, AND N. J. TRASK (1968) Nappes and gneiss domes in west-central New England. In E-an Zen, W. S. White, J. B. Hadley, and J. B. Thompson, Jr., Eds., *Studies of Appalachian Geology, Northern and Maritime*. John Wiley and Sons, New York, p. 203-218.

*Manuscript received, November 8, 1973; accepted for publication, February 26, 1974.*