

Halogen heterogeneity in the subcontinental lithospheric mantle revealed by I/Br ratios in kimberlites and their mantle xenoliths from South Africa, Greenland, China, Siberia, Canada, and Brazil

CHIAKI TOYAMA^{1,2}, HIROCHIKA SUMINO^{3,*·†}, NOBUAKI OKABE^{2,4}, AKIRA ISHIKAWA⁵, JUNJI YAMAMOTO⁶, ICHIRO KANEOKA⁷, AND YASUYUKI MURAMATSU²

¹Institute of Geology and Geoinformation, Geological Survey of Japan, The National Institute of Advanced Industrial Science and Technology (AIST), Ibaraki 237-0061, Japan

²Department of Chemistry, Gakushuin University, Tokyo 171-8588, Japan

³Department of General Systems Studies, Graduate School of Arts and Sciences, The University of Tokyo, Tokyo 153-8902, Japan

⁴Health and Environmental Risk Division, National Institute for Environmental Studies, Tsukuba 305-8506, Japan

⁵Department of Earth and Planetary Sciences, School of Science, Tokyo Institute of Technology, Tokyo 152-8851, Japan

⁶Department of Earth and Planetary Sciences, Graduate School of Science, Kyushu University, Fukuoka 819-0395, Japan

⁷Earthquake Research Institute, The University of Tokyo, Tokyo 113-0032, Japan

ABSTRACT

To investigate halogen heterogeneity in the subcontinental lithospheric mantle (SCLM), we measured the concentrations of Cl, Br, and I in kimberlites and their mantle xenoliths from South Africa, Greenland, China, Siberia, Canada, and Brazil. The samples can be classified into two groups based on halogen ratios: a high-I/Br group (South Africa, Greenland, Brazil, and Canada) and a low-I/Br group (China and Siberia). The halogen compositions were examined with the indices of crustal contamination using Sr and Nd isotopes and incompatible trace elements. The results indicate that the difference between the two groups was not due to different degrees of crustal contamination but from the contributions of different mantle sources. The low-I/Br group has a similar halogen composition to seawater-influenced materials such as fluids in altered oceanic basalts and eclogites and fluids associated with halite precipitation from seawater. We conclude that the halogens of the high-I/Br group are most likely derived from a SCLM source metasomatized by a fluid derived from subducted serpentinite, whereas those of the low-I/Br group are derived from a SCLM source metasomatized by a fluid derived from seawater-altered oceanic crust. The SCLM beneath Siberia and China could be an important reservoir of subducted, seawater-derived halogens, while such role of SCLM beneath South Africa, Greenland, Canada, and Brazil seems limited.

Keywords: Halogens, kimberlite, subcontinental lithospheric mantle, subduction, metasomatism; Halogens in Planetary Systems