

SUPPORTING INFORMATION (SI)

Calorimetric study of skutterudite (CoAs_{2.92}) and heazlewoodite (Ni₃S₂)

JURAJ MAJZLAN^{1,*}, STEFAN KIEFER¹, KRISTINA LILOVA², TAMILARASAN SUBRAMANI²,
ALEXANDRA NAVROTSKY², MAREK TUHÝ^{3,4}, ANNA VYMAZALOVÁ³, DMITRIY A. CHAREEV^{5,6,7},
EDGAR DACHS⁸, ARTUR BENISEK⁸

¹ Institute of Geosciences, Friedrich-Schiller University, Burgweg 11, 07749 Jena, Germany

² School of Molecular Sciences and Center for Materials of the Universe, Arizona State University, Tempe, Arizona 85287, USA

³ Czech Geological Survey, Geologická 6, 152 00 Prague 5, Czech Republic

⁴ Institute of Geochemistry, Mineralogy and Mineral Resources, Faculty of Science, Charles University, Albertov 6, 128 00 Prague

⁵ Institute of Experimental Mineralogy (IEM RAS), 142432 Chernogolovka, Moscow Region, Russia

⁶ Ural Federal University, Ekaterinburg 620002, Russia

⁷ Dubna State University, Dubna 141982 Russia

⁸ Department of Chemistry and Physics of Materials, University of Salzburg, Jakob-Haringer-Strasse 2a, 5020 Salzburg, Austria

*corresponding author: email: Juraj.Majzlan@uni-jena.de
telephone: +49-3641-948700
fax: +49-3641-948602
ORCID: 0000-0002-9151-4394

Number of figures: 2

Number of tables: 3

Number of pagers in SI (including the cover page): 4

Table S1. Equilibrium constants for selected reactions considered in this work. The values were calculated by SUPCRTBL (Zimmer et al. 2016). P_{sat} is the saturation H_2O pressure along the water-water vapor curve. For these calculations, the data for heazlewoodite and skutterudite were taken from this work, data for all other phases and species in reactions (11) through (14) from SUPCRTBL.

T (°C)	P_{sat} (bar)	$\log K_{11}$	$\log K_{12}$	$\log K_{13}$	$\log K_{14}$
0.01	1	44.176	29.344	207.802	179.758
25	1	39.929	25.954	190.208	163.51
50	1	36.353	22.91	175.067	149.407
75	1	33.31	20.185	161.915	137.059
100	1.013	30.69	17.725	150.377	126.137
125	2.32	28.405	15.477	140.162	116.385
150	4.757	26.386	13.394	131.04	107.594
175	8.918	24.579	11.438	122.832	99.6
200	15.536	22.942	9.576	115.393	92.267
225	25.479	21.439	7.773	108.606	85.479
250	39.736	20.032	5.987	102.371	79.128
275	59.431	18.693	4.176	96.6	73.107
300	85.838	17.416	2.309	91.223	67.305
325	120.458	16.242	0.371	86.175	61.573
350	165.211	15.304	-1.695	81.37	55.533

Table S2. Adjustable parameters A_1 - A_5 for the equation (15) for the chemical reactions (11) through (14). The fitted $\log K$ values are listed in Table S1.

	$\log K_{11}$	$\log K_{12}$	$\log K_{13}$	$\log K_{14}$
A_1	3.9819E+00	-3.1778E+03	-1.2690E+03	-5.7571E+03
A_2	-2.1705E-02	-4.8008E-01	-1.9782E-01	-8.3650E-01
A_3	6.3566E+03	1.9026E+05	1.3555E+05	3.8959E+05
A_4	4.3749E+00	1.1444E+03	4.5449E+02	2.0625E+03
A_5	9.1092E+05	-1.0930E+07	-5.4261E+06	-2.1345E+07

Table S3. Adjustable parameters a-f for the equation (16) for the chemical reactions (11) through (14). The fitted $\log K$ values are listed in Table S1.

	$\log K_{11}$	$\log K_{12}$	$\log K_{13}$	$\log K_{14}$
a	3.9951E+01	2.5947E+01	1.9020E+02	1.6346E+02
b	-2.4488E+00	1.4112E+00	7.3071E-01	6.8141E+00
c	7.1222E-04	-5.5498E-04	-2.7248E-04	-2.2451E-03
d	4.1998E+05	-1.3204E+05	-2.2692E+04	-9.1423E+05
e	-1.9768E+07	5.1839E+06	2.4850E+06	4.3839E+07
f	1.5208E+03	-6.8653E+02	-3.8370E+02	-3.8921E+03

References

- Zimmer, K., Zhang, Y.L., Lu, P., Chen, Y.Y., Zhang, G.R., Dalkilic, M., and Zhu, C. (2016) SUPCRTBL: A revised and extended thermodynamic dataset and software package of SUPCRT92. *Computer and Geosciences*, 90, 97-111. <https://doi.org/10.1016/j.cageo.2016.02.013>




Fig. S1. Powder X-ray diffraction pattern of synthetic heazlewoodite used in this work.




Fig. S2. Powder X-ray diffraction pattern of synthetic skutterudite used in this work.