

## **Analytical techniques for volatiles: A case study using intermediate (andesitic) glasses**

**P.L. KING,<sup>1,\*</sup> T.W. VENNEMANN,<sup>2</sup> J.R. HOLLOWAY,<sup>1,3</sup> R.L. HERVIG,<sup>1,4</sup> J.B. LOWENSTERN,<sup>5</sup> AND J.F. FORNERIS<sup>1</sup>**

<sup>1</sup>Geology Department, Arizona State University, Tempe, Arizona 85287, U.S.A.

<sup>2</sup>Institut Geochemie, Wilhelmstr. 56, D-72076, Tübingen, Germany

<sup>3</sup>Chemistry Department, Arizona State University, Tempe, Arizona 85287, U.S.A.

<sup>4</sup>Center for Solid State Science, Arizona State University, Tempe, Arizona 85287, U.S.A.

<sup>5</sup>U.S. Geological Survey, MS 910, 345 Middlefield Road, Menlo Park, California 94025, U.S.A.

### **ABSTRACT**

Small-scale analyses of volatiles in minerals and glasses provide information on how volatiles influence high-temperature geologic processes and low-temperature alteration processes. Four techniques for determining the C-O-H volatile contents of andesitic glasses are compared: manometry, secondary ion mass spectrometry, micro-Fourier transform infrared spectroscopy, and a technique where the H<sub>2</sub>O content is calculated using the difference between electron microprobe analysis totals and 100% sum. We present a method to determine the H content of a wide range of glass and mineral compositions using secondary ion mass spectrometry and a model for calibration factors. The extinction coefficients for H-O volatile contents in intermediate composition synthetic glasses are determined, and it is demonstrated that C-O speciation changes as total H<sub>2</sub>O content increases, with molecular CO<sub>2</sub> decreasing, CO<sub>3</sub><sup>2-</sup> increasing, and carbonate peak splitting increasing. For glasses with low H<sub>2</sub>O content and oxy-substituted minerals, the methods of choice for volatile analysis are secondary ion mass spectrometry or micro-Fourier transform infrared spectroscopy.