

## **Ni-phylosilicates (garnierites) from the Falcondo Ni-laterite deposit (Dominican Republic): Mineralogy, nanotextures, and formation mechanisms by HRTEM and AEM**

**CRISTINA VILLANOVA-DE-BENAVENT<sup>1,\*</sup>, FERNANDO NIETO<sup>2</sup>, CECILIA VITI<sup>3</sup>, JOAQUÍN A. PROENZA<sup>1</sup>, SALVADOR GALÍ<sup>1</sup>, AND JOSEP ROQUÉ-ROSELL<sup>4</sup>**

<sup>1</sup>Departament de Cristal·lografia, Mineralogia i Dipòsits Minerals, Facultat de Geologia, Universitat de Barcelona (UB), Martí i Franquès s/n, 08028 Barcelona, Spain

<sup>2</sup>Departamento de Mineralogía y Petrología and IACT, Universidad de Granada, CSIC, Av. Fuentenueva, 18071 Granada, Spain

<sup>3</sup>Dipartimento di Scienze Fisiche, della Terra e dell'Ambiente, Università degli Studi di Siena, Via Laterina 8, 53100 Siena, Italy

<sup>4</sup>Advanced Light Source, Lawrence Berkeley National Laboratory, One Cyclotron Road, MS 15R0317, Berkeley, California 94720, U.S.A.

### **ABSTRACT**

Ni-bearing magnesium phyllosilicates (garnierites) are significant Ni ores in Ni-laterites worldwide. The present paper reports a detailed TEM investigation of garnierites from the Falcondo Ni-laterite deposit (Dominican Republic). Different types of garnierites have been recognized, usually consisting of mixtures between serpentine and talc-like phases that display a wide range of textures at the nanometer scale. In particular, chrysotile tubes, polygonal serpentine, and lizardite lamellae are intergrown with less crystalline, talc-like lamellae. Samples consisting uniquely of talc-like and of sepiolite-falcondoite were also observed, occurring as distinctive thin lamellae and long ribbon-shaped fibers, respectively. HRTEM imaging indicates that serpentine is replaced by the talc-like phase, whereas TEM-AEM data show preferential concentration of Ni in the talc-like phase. We suggest, therefore, that the crystallization of Ni-bearing phyllosilicates is associated with an increase in the silica activity of the system, promoting the replacement of the Ni-poor serpentine by the Ni-enriched talc-like phase. These results have interesting implications in material science, as garnierites are natural analogs of Ni-bearing phyllosilicate-supported synthetic catalysts. Finally, SAED and HRTEM suggest that the Ni-bearing talc-like phase corresponds to a variety of talc with extra water, showing larger  $d_{001}$  than talc (i.e., 9.2–9.7 Å), described as “kerolite”-“pimelite” in clay mineral literature.

**Keywords:** Ni-laterites, garnierites, chrysotile, polygonal serpentine, lizardite, “kerolite”-“pimelite”, sepiolite-falcondoite, HRTEM