

Chromium-bearing phases in the Earth's mantle: Evidence from experiments in the Mg₂SiO₄–MgCr₂O₄ system at 10–24 GPa and 1600 °C

EKATERINA A. SIROTKINA^{1,2,3,*}, ANDREY V. BOBROV^{1,2,3}, LUCA BINDI^{4,5}, AND TETSUO IRIFUNE^{6,7}

¹Vernadsky Institute of Geochemistry and Analytical Chemistry of Russian Academy of Sciences, Moscow 119991, Russia

²Department of Petrology, Geological Faculty, Moscow State University, Moscow 119991, Russia

³Institute of Experimental Mineralogy of Russian Academy of Sciences, Chernogolovka 142432, Russia

⁴Dipartimento di Scienze della Terra, Università di Firenze, Via La Pira 4, 50121 Florence, Italy

⁵CNR, Istituto di Geoscienze e Georisorse, sezione di Firenze, Via La Pira 4, 50121 Florence, Italy

⁶Geodynamics Research Center, Ehime University, Matsuyama 790-8577, Japan

⁷Earth-Life Science Institute, Tokyo Institute of Technology, Tokyo 152-8550, Japan

ABSTRACT

Phase relations in the system Mg₂SiO₄–MgCr₂O₄ were studied at 10–24 GPa and 1600 °C using a high-pressure Kawai-type multi-anvil apparatus. We investigated the full range of starting compositions for the forsterite-magnesiochromite system to derive a *P*–*X* phase diagram and synthesize chromium-bearing phases, such as garnet, wadsleyite, ringwoodite, and bridgmanite of a wide compositional range. Samples synthesized at 10 GPa contain olivine with small chromium content and magnesiochromite. Mg₂SiO₄ wadsleyite is characterized by the pressure-dependent higher chromium solubility (up to 7.4 wt% Cr₂O₃). The maximal solubility of chromium in ringwoodite in the studied system (~18.5 wt% Cr₂O₃) was detected at *P* = 23 GPa, which is close to the upper boundary of the ringwoodite stability. Addition of chromium to the system moves the boundaries of olivine/wadsleyite and wadsleyite/ringwoodite phase transformations to lower pressures. Our experiments simulate Cr-rich phase assemblages found as inclusions in diamonds, mantle xenoliths, and UHP podiform chromitites.

Keywords: Magnesiochromite, forsterite, olivine, wadsleyite, ringwoodite, knorringite, majorite, bridgmanite, mantle, high-*P*-*T* experiments, phase relations