

Hartite from Bílina

VLADIMÍR BOUŠKA,^{1,*} IVANA ČÍSAŘOVÁ,¹ ROMAN SKÁLA,² ZDENĚK DVOŘÁK,³ JAROSLAV ZELINKA,¹ and KAREL ŽÁK²

¹Faculty of Science, Charles University, Albertov 6, CZ-128 43 Prague, Czech Republic

²Czech Geological Survey, Klárov 3/131, CZ-118 21 Prague 1, Czech Republic

³The Bílina Mines, The North Bohemian Mines Co., 5.kvetna str., CZ-418 29 Bílina, Czech Republic

ABSTRACT

In the Bílina Mine (Czech Republic), a very chemically pure material corresponding to hartite was found as a product of distillation into the pelosiderite nodule during coalification of a lignite seam in the Miocene or later. The amount of the material available allowed characterizing this natural mineral equivalent of the hydrocarbon (+)-phylocladane [16 α (H)-phylocladane] thoroughly.

Elementary analysis of hartite from Bílina gave C = 87.45 wt% and H = 12.55 wt%, which corresponds to the formula C₂₀H₃₄. The IR spectrum demonstrated the presence of gem-dimethyl groups (doublet δ_s CH₃ at 1369 and 1383 cm⁻¹). Other bands indicated the presence of methyl and methylene groups. NMR studies showed that the hydrocarbon (hartite) molecule contains four CH₃ groups, nine CH₂ groups, four CH groups, and three quaternary carbon atoms. The sample exhibits a well-defined melting point between 71 and 71.5 °C. Measurements of optical rotation gave $[\alpha]_D = +24.2^\circ$ (CHCl₃, $c = 0.7$). Lattice parameters from single-crystal study measured at 293(2) K were $a = 11.407$ (1), $b = 20.952$ (2), $c = 7.4060$ (8) Å, $\alpha = 93.941$ (9)°, $\beta = 100.750$ (8)°, and $\gamma = 80.499$ (9)°. The crystal system is triclinic with space group *Pl*, $Z = 4$, $V = 1713.8$ (3) Å³. Densities are $D_m = 1.04$ g/cm³ and $D_x = 1.064$ g/cm³, respectively. Isotopic composition of C in hartite from Bílina gave $\delta^{13}\text{C}_{(\text{PDB})} = -24.4(1)$ ‰, a common value for coal and organic sedimentary material derived from higher plants.