1	Revision 1-shortened version
2	Magma oxygen fugacity of mafic-ultramafic intrusions in
3	convergent margin settings: insights for the role of magma
4	oxidation states on magmatic Ni-Cu sulfide mineralization
5	
6	
7	Yonghua Cao ^{1, 2} , Christina Yan Wang ^{1, 2} *, Bo Wei ^{1, 2}
8	¹ CAS Key Laboratory of Mineralogy and Metallogeny, Guangzhou Institute of
9	Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China
10	² Guangdong Provincial Key Laboratory of Mineral Physics and Materials, Guangzhou
11	510640, China
12	
13	
14	
15	
16	
17	Corresponding author: Dr. C.Y. Wang (wang_yan@gig.ac.cn)

1

18

Abstract

Oxygen fugacities (fO_2) of mantle-derived mafic magmas have important controls on 19 20 the sulfur status and solubility of the magmas, which are key factors to the formation of 21 magmatic Ni-Cu sulfide deposits, particularly those in convergent margin settings. In 22 order to investigate the fO_2 of mafic magmas related to Ni-Cu sulfide deposits in 23 convergent margin settings, we obtained the magma fO_2 of a number of Ni-Cu sulfide-24 bearing mafic-ultramafic intrusions in the central Asian orogenic belt (CAOB), North 25 China, based on the olivine-spinel oxygen barometer and the modeling of V partitioning 26 between olivine and melt. We also calculated the mantle fO_2 on the basis of V/Sc ratios 27 of primary magmas of these intrusions.

Ni-Cu sulfide-bearing mafic-ultramafic intrusions in the CAOB include arc-related 28 29 Silurian-Carboniferous ones and post-collisional Permian-Triassic ones. Arc-related intrusions formed before the closure of the paleo-Asian ocean and include the Jinbulake, 30 31 Heishan, Kuwei and Erbutu intrusions. Post-collisional intrusions were emplaced in 32 extensional settings after the closure of the paleo-Asian ocean and include the Kalatongke, Baixintan, Huangshandong, Huangshan, Poyi, Poshi, Tulaergen and Hongqiling No.7 33 intrusions. It is clear that the magma fO_2 values of all these intrusions in both settings 34 range mostly from FMQ+0.5 to FMQ+3 and are generally elevated with the fractionation 35 36 of magmas, much higher than that of MORBs (FMQ-1 to FMQ+0.5). However, the 37 mantle fO_2 values of these intrusions vary from ~FMQ to ~FMQ+1.0, just slightly higher 38 than that of MORBs (\leq FMQ). This slight difference is interpreted as the intrusions in the 39 CAOB may have been derived from the metasomatized mantle wedges where only minor 40 slab-derived, oxidized components were involved. Therefore, the high magma fO_2 values

41	of most Ni-Cu sulfide-bearing mafic-ultramafic intrusions in the CAOB were attributed
42	to the fractionation of magmas derived from the slightly oxidized metasomatized mantle.
43	In addition, the intrusions that host economic Ni-Cu sulfide deposits in the CAOB usually
44	have magma fO_2 of >FMQ+1.0 and sulfides with mantle-like $\delta^{34}S$ values (-1.0 to +1.1‰)
45	indicating that the oxidized mafic magmas may be able to dissolve enough mantle-
46	derived sulfur to form economic Ni-Cu sulfide deposits. Oxidized mafic magmas derived
47	from metasomatized mantle sources may be an important feature of major orogenic belts.
48	Keywords: Mafic-ultramafic intrusion; Magmatic Ni-Cu sulfide mineralization; Magma
49	oxygen fugacity; Central Asian orogenic belt; Convergent margin setting

- 50
- 51

INTRODUCTION

The oxygen fugacity (fO_2) of mantle-derived mafic magmas is controlled by 52 equilibria of Fe³⁺-Fe²⁺ and S²⁻-S⁶⁺ (Kress and Carmichael, 1991; Jugo et al., 2005), and 53 54 can be quantified as $\triangle \log fO_2$ relative to mineral assemblage buffers. The fO_2 values of 55 mafic magmas are considered to be closely related to geodynamic settings, but how they differ in different settings is still a matter of debate. In general, having $Fe^{3+}/\Sigma Fe$ and 56 $S^{6+}/\Sigma S$ higher than the mid-ocean ridge basalts (MORBs), arc and back-arc basalts may 57 58 have formed from relatively oxidized magmas (Wood et al., 1990; Nilsson and Peach, 59 1993; Jugo et al., 2010; Brounce et al., 2017). It has been demonstrated that arc and backarc basalts were derived from metasomatized mantle wedges that have been oxidized to 60 61 variable degrees (Debret et al., 2016; Rielli et al., 2017; Bénard et al., 2018). It is also 62 known that the metasomatized mantle beneath subduction zones has fO_2 similar to the mantle beneath the mid-ocean ridges, and it is the fractionation of metasomatized mantle-63

derived magmas or the interaction of hydrated magmas with ambient mantle that elevated the magma fO_2 (Lee et al., 2005, 2010; Dauphas et al., 2010; Tollan and Hermann, 2019; Li et al., 2020).

67 Magmatic Ni-Cu sulfide deposits are traditionally thought to be related to the mafic magmatism induced by either mantle plumes or rifting within intraplate settings (Naldrett, 68 2004). However, mafic-ultramafic intrusions in convergent margin settings have become 69 70 targets for prospecting economic Ni-Cu sulfide deposits in recent years (Maier et al., 71 2008; Thakurta et al., 2008; Tomkins et al., 2012; Manor et al., 2016; Song et al., 2016). 72 The mantle sources of such intrusions in are generally considered to be metasomatized by 73 slab-derived fluids/melts (Manor et al., 2016; Song et al., 2016). The mafic magmas derived from the metasomatized mantle can be highly hydrated and oxidized with fO_2 74 75 being up to FMO+6 (FMO means fayalite-magnetite-quartz oxygen buffer) (Kelley and Cottrell, 2009; Kelley et al., 2010; Gaillard et al., 2015). For example, the magma fO_2 of 76 77 the Alaskan-type Duke intrusion in USA and the Turnagain and Mascot Ni-Cu sulfide-78 bearing mafic-ultramafic intrusions in Spain are calculated to be >FMQ+2 (Thakurta et al., 2008; Manor et al., 2016). The central Asian orogenic belt (CAOB) is one of the 79 80 largest accretionary orogens in the world, resulted from large-scaled subduction and accretion of juvenile materials from Neoproterozoic to Paleozoic (Sengör et al., 1993; 81 82 Xiao et al., 2004a, b, 2009; Jahn et al., 2004). A preliminary study on the oxidation states 83 of a few Ni-Cu sulfide-bearing mafic-ultramafic intrusions in the CAOB indicates that 84 magma fO_2 values vary from FMQ+0.3 to FMQ+2.6, much higher than that of MORBs 85 (Cao et al., 2019).

86 Experimental results indicate that the sulfur solubility of highly oxidized mafic magmas can be as high as 1.4 wt.% with sulfur being dominantly as sulfate species (S^{6+}) 87 88 (Jugo et al., 2005; Jugo, 2009), significantly higher than that of reduced mafic magmas with dominantly S²⁻ phases (Jugo et al., 2010; Cottrell and Kelley, 2011). Therefore, the 89 oxidized mantle source or highly oxidized, hydrated mafic magmas may be more 90 91 favorable for the magmatic Ni-Cu sulfide deposits in convergent margin settings (Jenner 92 et al., 2010; Tomkins et al., 2012; Cao et al., 2019; Wei et al., 2019). However, the linkage between magma fO_2 of mafic-ultramafic intrusions and Ni-Cu sulfide 93 94 mineralization is not well understood. Three important issues that should be answered: (1) 95 if the mantle sources of the mafic-ultramafic intrusions in convergent margin settings 96 have remarkably high fO_2 relative to those in intraplate settings? (2) if not, what triggers 97 high magma fO_2 of the mafic-ultramatic intrusions in convergent margin settings? and (3) what is the favorable magma fO_2 for the Ni-Cu sulfide mineralization in convergent 98 99 margin settings?

100 A number of Paleozoic mafic-ultramafic intrusions in the CAOB host Ni-Cu sulfide deposits with variable Ni grades and ore reserves, making up a ~4000-km-long Ni-Cu 101 102 sulfide mineralization belt in North China. These intrusions were dated to be Devonian to 103 Triassic in ages, some of which were emplaced in the subduction stage predating the 104 closure of the paleo-Asian ocean, whereas others in the post-subduction, extensional 105 stage after the closure of the paleo-Asian ocean (e.g., Yang and Zhou, 2009; Qin et al., 2011; Li et al., 2012; Yang et al., 2012; Peng et al., 2013; Li et al., 2015). These 106 107 intrusions are ideal to unravel the correlation between magma fO_2 and Ni-Cu sulfide 108 mineralization in a convergent margin setting. In this study, we estimated the mantle and

magma fO_2 of representative mafic-ultramafic intrusions in the CAOB that were emplaced in different ages and host variable degrees of Ni-Cu sulfide mineralization. The results indicate that most intrusions have magma fO_2 much higher than that of MORBs despite the similarity in their mantle fO_2 . Such a feature can be further examined for the Ni-Cu sulfide-bearing mafic-ultramafic intrusions in convergent margin settings elsewhere.

- 115
- 116

GEOLOGICAL BACKGROUND

The central Asian orogenic belt is bounded by the Siberian Craton to the north and the Tarim Craton and North China Craton to the south (Fig. 1a). The belt extends for more than 7000 km from the Pacific ocean to the Eastern Europe, making up one of the largest accretionary orogenic belts on Earth. It formed due to the closure of the paleo-Asian ocean in Paleozoic and comprises numerous fragments of Precambrian microcontinents, Paleozoic island arcs, ophiolite suites, successions of volcanic rocks (Windley et al., 2007; Xiao et al., 2009).

The CAOB in China part is subdivided into the western and eastern segments (Zhou 124 125 and Wilde, 2013) (Fig. 1b). The western segment is further divided into five belts, from north to south (Fig. 1c), including: 1) the Altay orogenic belt that is bounded by the 126 127 Sayan belt to the north and by the Ulungar fault and Junggar block to the south (Sengör et 128 al., 1993; Windley et al., 2002; Xiao et al., 2009), 2) the North Tianshan orogenic belt 129 between the Junggar block to the north and the Aqikkuduk fault to the south (Zhou et al., 130 2004; Qin et al., 2011; Gao et al., 2012), 3) the Central Tianshan orogenic belt between 131 the Aqikkuduk fault to the north and the Kawabulak fault to the south (Song et al., 2013),

4) the South Tianshan orogenic belt between the Kawabulak fault to the north and the
Tarim Craton to the south (Yang and Zhou, 2009), and 5) the Beishan fold belt along the
northeastern margin of the Tarim Craton (Xu et al., 2016). The eastern segment refers to
the Xing'an-Mongolia orogenic belt in the Inner Mongolia and NE China (Zhang et al.,
2015), which consists mainly of, from north to south, the Erguna massif, Xing'an massif,
Songnen-Zhangguangcai range massif, and a continental margin accretionary belt (Wu et al., 2007) (Fig. 1d).

Numerous mafic-ultramafic intrusions that contain Ni-Cu sulfide mineralization occur in the CAOB. They were emplaced mainly in two periods, one from Silurian to Carboniferous and the other from Permian to Triassic (*e.g.*, Yang and Zhou, 2009; Xie et al., 2012; Hao et al., 2014; Mao et al., 2016).

143 Silurian to Carboniferous mafic-ultramafic intrusions

Silurian to Carboniferous mafic-ultramafic intrusions are mainly distributed in the 144 145 western segment of the CAOB and host small- to medium-sized Ni-Cu sulfide deposits 146 (Fig. 1b). As the paleo-Asian ocean was not yet closed until Permian in the western 147 segment (Han et al., 2007; Xiao et al., 2009), these intrusions are considered to be arc-148 related (Yang and Zhou, 2009; Xie et al., 2012; Yang et al., 2012). Representative intrusions include the Jinbulake intrusion (ca. 430 Ma) in the central Tianshan belt (Yang 149 and Zhou, 2009; Yang et al., 2012), the Kuwei intrusion (ca. 398 Ma) in the Altay belt 150 151 (Li et al., 2015), and the Heishan intrusion (ca. 356 to 367 Ma) in the Beishan belt (Xie et 152 al., 2012).

The parental magmas of these intrusions are tholeiitic (*e.g.*, Zhou et al., 2004; Yang and Zhou, 2009; Tang et al., 2012; Xia et al., 2013; Song et al., 2013). Rocks of these

intrusions have positive $\varepsilon_{Nd}(t)$ (+0.4 to +4) and initial Sr⁸⁷/Sr⁸⁶ ranging from 0.704 to 0.709 (Yang and Zhou, 2009; Xie et al., 2012; Yang et al., 2012). They show depleted Nb and Ta relative to large ion lithophile elements (LILE) and light rare earth elements (LREE) on the primitive mantle-normalized trace element patterns (Fig. 2a-d), consistent with an arc-like affinity. These features were interpreted as magma generation from the depleted mantle that had been metasomatized by slab-derived fluids/melts (Yang and Zhou, 2009; Xie et al., 2012; Yang et al., 2012).

The Erbutu intrusion in the eastern segment of the CAOB is an outlier. Although it is dated to be 294.2 ± 2.7 Ma, it is considered to be an arc-hosted intrusion (Peng et al., 2013). The intrusion hosts a small-sized Ni-Cu sulfide deposit and the parental magma is boninitic (Peng et al., 2013). The intrusion is mainly composed of olivine-bearing orthopyroxenite with mineral modes quite similar to those formed from boninitic magma (Peng et al., 2013). The rocks have LREE and LILE (*e.g.*, Ba and Rb) more enriched than those of the Jinbulake and Heishan intrusions (Fig. 2e, f).

169 **Permian to Triassic mafic-ultramafic intrusions**

170 Permian to Triassic mafic-ultramafic intrusions in the CAOB host a number of 171 economic Ni-Cu sulfide deposits, including the Kalatongke intrusion (290-282 Ma) in the 172 Altay belt (Song and Li, 2009; Zhang et al., 2009; Gao et al., 2012), the Huangshandong 173 and Huangshanxi intrusions (274-283 Ma) in the Huangshan-Jingerquan mineralized belt in the North Tianshan belt (Qin et al., 2011; Sun et al., 2013), the Tulaergen intrusion 174 (265±9.2 Ma) in the Kanggur-Huangshan shear zone in the North Tianshan belt (Zhao et 175 176 al., 2017), the Poyi and Poshi intrusions (270-277 Ma) in the Beishan belt (Xue et al., 177 2016), and the Hongqiling No.7 and Piaohechuan No.4 intrusions (ca. 210-230 Ma) in the

Xing'an-Mongolia belt (Wei et al., 2013, 2015) (Fig. 1b). In addition, many other
intrusions in this period host potential Ni-Cu sulfide mineralization, including the
Huangshannan (278±2 Ma) and Baixintan intrusions (286±3 Ma) in the North Tianshan
belt (Mao et al., 2016; Feng et al., 2017), the Luodong intrusion (260-290 Ma) in the
Beishan belt (Su et al., 2015), and the Hongqiling No.1, 2, 3, 9, 32 and 33 intrusions (ca.
210-230 Ma) in the Xing'an-Mongolia belt (Hao et al., 2014).

These intrusions are considered to have formed in post-subduction, extensional 184 settings after the closure of the paleo-Asian ocean (e.g., Jiang et al., 2009; Li et al., 2012; 185 186 Sun et al., 2013; Wei et al., 2013, 2015; Mao et al., 2014, 2015). The rocks of these intrusions show arc-like trace element patterns (Fig. 3a-d), which are attributed to the 187 188 derivation from the metasomatized, depleted mantle (Xie et al., 2012; Li et al., 2012; 189 Mao et al., 2014; Deng et al., 2015). However, the rocks of the Luodong intrusion have 190 MORB-like, LREE-depleted trace element patterns (Fig. 3e, f), which may have been 191 derived from the weakly metasomatized mantle (Su et al., 2015).

192

193 INTRUSIONS AND SAMPLES CHOSEN FOR OXYGEN FUGACITY

194

CALCULATION

A prerequisite to use the olivine-spinel oxygen barometer is to obtain the compositions of equilibrated olivine-spinel pair in rocks (Ballhaus et al., 1991). The mafic-ultramafic intrusions in the CAOB that have rocks containing olivine-spinel pair include Silurian to Carboniferous Jinbulake, Heishan and Erbutu intrusions, and Permian to Triassic Baixintan, Huangshannan, Huangshandong, Huangshanxi, Poyi, Luodong, Tulaergen, Hongqiling No.1 and No. 2 intrusions. In this study, we calculated the magma

201	and mantle fO_2 values of the Jinbulake, Heishan, Erbutu, Baixintan, Huangshannan,					
202	Luodong, and Tulaergen intrusions. Together with the magma and/or mantle fO_2 values					
203	of the Huangshandong, Huangshanxi, Poyi and Hongqiling No.1 and No. 2 intrusions that					
204	were obtained in our earlier studies (Cao et al., 2019; Wei et al., 2019), an integrated					
205	framework of the magma and mantle fO_2 of the Ni-Cu sulfide-bearing mafic-ultramafic					
206	intrusions in the CAOB can be outlined. The results in this study are compared with the					
207	magma fO_2 values of the picrite in the Dali area, SW China, which is part of the					
208	Emeishan large igneous province (LIP) that formed within an intraplate setting. The					
209	petrography of the selected mafic-ultramafic intrusions in the CAOB and the Dali picrite					
210	in the Emeishan LIP were described in Supplementary Information.					
211						
212	ANALYTICAL RESULTS					
213	Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB					
213 214	Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB The compositions of the olivine-spinel pairs in the rocks of the selected mafic-					
213214215	Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB The compositions of the olivine-spinel pairs in the rocks of the selected mafic- ultramafic intrusions in the CAOB were analyzed in this study. Analytical methods were					
213214215216	Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB The compositions of the olivine-spinel pairs in the rocks of the selected mafic- ultramafic intrusions in the CAOB were analyzed in this study. Analytical methods were described in Supplementary Information. The results of the olivine-spinel pairs are					
 213 214 215 216 217 	Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB The compositions of the olivine-spinel pairs in the rocks of the selected mafic- ultramafic intrusions in the CAOB were analyzed in this study. Analytical methods were described in Supplementary Information. The results of the olivine-spinel pairs are described in Supplementary Information and the data are listed in Table S1.					
 213 214 215 216 217 218 	Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB The compositions of the olivine-spinel pairs in the rocks of the selected mafic- ultramafic intrusions in the CAOB were analyzed in this study. Analytical methods were described in Supplementary Information. The results of the olivine-spinel pairs are described in Supplementary Information and the data are listed in Table S1.					
 213 214 215 216 217 218 219 	Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB The compositions of the olivine-spinel pairs in the rocks of the selected mafic- ultramafic intrusions in the CAOB were analyzed in this study. Analytical methods were described in Supplementary Information. The results of the olivine-spinel pairs are described in Supplementary Information and the data are listed in Table S1. A summary of spinel compositions					
 213 214 215 216 217 218 219 220 	 Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB The compositions of the olivine-spinel pairs in the rocks of the selected mafic-ultramafic intrusions in the CAOB were analyzed in this study. Analytical methods were described in Supplementary Information. The results of the olivine-spinel pairs are described in Supplementary Information and the data are listed in Table S1. A summary of spinel compositions The spinel grains from the mafic-ultramafic intrusions in either arc or post-subduction, 					
 213 214 215 216 217 218 219 220 221 	Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB The compositions of the olivine-spinel pairs in the rocks of the selected mafic- ultramafic intrusions in the CAOB were analyzed in this study. Analytical methods were described in Supplementary Information. The results of the olivine-spinel pairs are described in Supplementary Information and the data are listed in Table S1. A summary of spinel compositions The spinel grains from the mafic-ultramafic intrusions in either arc or post-subduction, extensional settings in the CAOB have highly variable Cr# and XFe ³⁺ . The grains from					
 213 214 215 216 217 218 219 220 221 222 	Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB The compositions of the olivine-spinel pairs in the rocks of the selected mafic- ultramafic intrusions in the CAOB were analyzed in this study. Analytical methods were described in Supplementary Information. The results of the olivine-spinel pairs are described in Supplementary Information and the data are listed in Table S1. A summary of spinel compositions The spinel grains from the mafic-ultramafic intrusions in either arc or post-subduction, extensional settings in the CAOB have highly variable Cr# and XFe ³⁺ . The grains from the Erbutu intrusion have the highest Cr# and the lowest XFe ³⁺ among the three arc-					
 213 214 215 216 217 218 219 220 221 222 223 	 Compositions of olivine-spinel pairs in mafic-ultramafic intrusions in the CAOB The compositions of the olivine-spinel pairs in the rocks of the selected mafic-ultramafic intrusions in the CAOB were analyzed in this study. Analytical methods were described in Supplementary Information. The results of the olivine-spinel pairs are described in Supplementary Information and the data are listed in Table S1. A summary of spinel compositions The spinel grains from the mafic-ultramafic intrusions in either arc or post-subduction, extensional settings in the CAOB have highly variable Cr# and XFe³⁺. The grains from the Erbutu intrusion have the highest Cr# and the lowest XFe³⁺ among the three arc-hosted intrusions (Fig. 4a, b). Among the intrusions in the post-subduction, extensional 					

224 settings, the spinel grains from the Baixintan, Huangshannan and Tulaergen intrusions have relatively restricted Cr# but highly variable XFe³⁺ relative to those from the 225 Luodong and Hongqiling No.1 and No.2 intrusions (Fig. 4a, b). In addition, the spinel 226 grains from the Luodong intrusion has similar Cr# but relatively low and restricted XFe³⁺ 227 228 compared to those from the Hongqiling No.1 and No.2 intrusions (Fig. 4a, b). The spinel grains from the Erbutu and Luodong intrusions are clustered on the plot of Mg# versus 229 XFe^{3+} , whereas the grains from each of other intrusions generally show a negative trend 230 of Mg# versus XFe³⁺on this plot (Fig. 4b). 231

The spinel grains in the Dali picrite overall have higher Mg# and Cr#, and lower XFe³⁺ than those from the intrusions in the CAOB (Fig. 4a, b). However, they have similar Cr# and XFe³⁺ to those from the Erbutu intrusion (Fig. 4a, b). They display a nearly horizontal trend on the plot of Mg# versus XFe³⁺ (Fig. 4b), which is in contrast to the negative correlation trend for the spinel from the intrusions in the CAOB on the plot.

237

238 S isotope compositions of sulfides in mafic-ultramafic intrusions in the CAOB

The method of *in situ* S isotope analysis for the sulfides (pyrrhotite, pentlandite and 239 chalcopyrite) in the rocks of the selected mafic-ultramafic intrusions in the CAOB is 240 241 described in Supplementary Information. The sulfides in the wehrlite of the Jinbulake intrusion have δ^{34} S ranging from +0.3 to +1.3% (Table 1). The sulfides in the lherzolite 242 of the Baixintan intrusion have δ^{34} S ranging from -0.7 to +1.2‰ (Table 1). The sulfides 243 in the lherzolite of the Tulaergen intrusion have δ^{34} S ranging from -0.2 to +0.8‰ (Table 244 1). Overall, the sulfides from the three intrusions have a restricted range of δ^{34} S from -0.7 245 to +1.3%. Likewise, the sulfides in the ores of three economic Ni-Cu sulfide deposits 246

hosted in the Permian-Triassic Kalatongke, Hongqiling No. 7 and Piaohechuan No. 4 intrusions in the CAOB have δ^{34} S ranging from -1.0 to +1.1‰ (Wei et al., 2019). All of these values are similar to the δ^{34} S of MORB-type mantle (-1.5 to +0.6‰, Labidi et al., 2013, 2014) (Fig. 5). In contrast, the sulfides from the rocks of the Erbutu intrusion have δ^{34} S ranging from +5.3 to +7.5‰ (Table 1), much higher than those from other intrusions in the CAOB (Fig. 5).

- 253
- 254

CALCULATION RESULTS OF OXYGEN FUGACITY

255 The oxygen fugacity of the mantle and mantle-derived mafic magmas can be calculated in four different ways, including: 1) measuring $Fe^{3+}/(Fe^{3+}+Fe^{2+})$ of basalts or 256 quenched basaltic glass (Kress and Carmichael, 1991; Kelley and Cottrell, 2009), 2) 257 258 quantifying the partition coefficients of redox-sensitive elements (e.g., V and Cr) in the differentiation of magma (Canil, 1997; Mallmann and O'Neill, 2009), 3) using oxygen 259 260 barometers based on the chemical equilibria between mineral pairs (e.g., olivine-spinel 261 pair) (Ballhaus et al., 1991), and 4) calculating the ratios of redox sensitive/insensitive elements (e.g., V/Sc, Fe^t/Zn) of primary magmas (Lee et al., 2005, 2010; Mallmann and 262 O'Neill, 2009). The fourth method is exclusively used to estimate the mantle oxygen 263 fugacity (Lee et al., 2005; Mallmann and O'Neill, 2009), however, the three others are 264 applicable to calculate the fO_2 of both mantle and mantle-derived magmas, depending on 265 that the examined objects are mantle xenoliths (e.g., Ionov and Wood, 1992), or 266 fractionated basalts/mafic-ultramafic intrusions (e.g., Cao et al., 2019). 267

268 Mantle fO_2

269 Given that the mantle xenolith that can be directly used to calculate the mantle fO_2 are unavailable in the CAOB, we constrained the mantle fO_2 based on the relationship 270 between the mantle fO_2 and the V/Sc ratios of primary magmas, an alternative method 271 272 proposed by Lee et al. (2005) and Mallmann and O'Neill (2009). Because V is sensitive to redox and Sc is not, the V/Sc ratio of primary magma is mainly governed by fO_2 273 274 during partial melting of a given mantle lithology (Lee et al., 2005; Mallmann and O'Neill, 2009), and is not affected by temperature and pressure (Canil and Fedortchouk, 275 276 2000; Li, 2018). In addition, the V/Sc ratio of basaltic magma is not sensitive to the 277 crystallization of olivine (Lee et al., 2005; Mallmann and O'Neill, 2009), the V/Sc ratio 278 of the melt in equilibrium with the most primitive olivine in a mafic-ultramafic intrusion can be taken as the ratio of primary magma, particularly if olivine is the only cumulus 279 280 phase. Therefore, we selected the samples from the Heishan, Huangshannan, Luodong, 281 Poyi and Hongqiling No.2 intrusions in the CAOB that contain high Fo olivine (Fo = 86282 to 90) as the only cumulus phase, the obtained V/Sc ratio of the melt in equilibrium with 283 the olivine is analog to the V/Sc ratio of the primary magma of the intrusion.

As olivine is the only cumulus phase in the rocks, the concentrations of V and Sc of the melt can be calculated using the mass balance equation (Godel et al., 2011):

$$C_{\rm WR}^{V,Sc} = F_{\rm Ol} \times C_{\rm Ol}^{V,Sc} + (1 - F_{\rm Ol}) \times C_{\rm Liq}^{V,Sc}$$
(1)

where $C_{WR}^{V, Sc}$ and $C_{Ol}^{V, Sc}$ is the concentrations of V and Sc in the bulk rock and cumulus olivine, respectively. The fraction of olivine (F_{Ol}) can be estimated in two ways; one is to analyze the back-scattered electron (BSE) images or scan thin sections of the samples, the other is to use the mass balance of whole-rock MgO and FeO contents combined with the olivine-liquid exchange coefficient (*Kd*) (Li and Ripley, 2011). In this study, we

292	integrated the two ways to obtain the F_{Ol} and then calculated the concentrations of V and
293	Sc in the melt $(C_{\text{Liq}}^{V, Sc})$ based on equation (1) (Table S2).

294 The V/Sc ratios of primary magmas would increase slightly with the degrees of 295 partial melting of the mantle at a given mantle fO_2 when it is \leq FMQ, but would decrease 296 significantly when it is >FMQ (Lee et al., 2005) (Fig. 6). Therefore, the degrees of partial 297 melting of the mantle should be considered when the V/Sc ratio of primary magma is 298 used to calculate mantle fO_2 . Mafic magmas in subduction zones are generally produced 299 by higher degrees of partial melting of the mantle (e.g., up to 15-20%, Kelley et al., 2006) than those in the mid-ocean ridges (~10%, Bottinga and Allegre, 1976). The degrees of 300 301 partial melting of the mantle are thus set to be 15 to 20% for the intrusions in the CAOB, 302 the obtained mantle fO_2 of the Heishan, Huangshannan, Luodong, Poyi and Hongqiling 303 No.2 intrusions is ~FMQ+1.0, ~FMQ, ~FMQ, ~FMQ+1.0 and ~FMQ+0.5, respectively 304 (Fig. 6).

In our previous study, the mantle fO_2 of the Poyi and Honggiling No.2 intrusions was 305 306 estimated to be FMQ+0.3 and FMQ+0.5, respectively, using the olivine-spinel oxygen 307 barometer (Cao et al., 2019). As the chemical data of the spinel from the Poyi intrusion in that study were collected from the literature and the $Fe^{3+}/\Sigma Fe$ of the spinel was not 308 corrected, the obtained mantle fO_2 was likely underestimated by ~0.6 log unit (Cao et al., 309 2019), so the mantle fO_2 of the Poyi intrusion could be ~FMQ+0.9. Therefore, the mantle 310 311 fO_2 of the Poyi and Hongqiling No.2 intrusions obtained by two different ways are quite 312 consistent with each other.

313 Magma *f*O₂

- The magma fO_2 of the mafic-ultramafic intrusions in the CAOB was acquired by two methods; one is based on the olivine-spinel oxygen barometer (Ballhaus et al., 1991), the other is based on V partitioning in olivine (Canil, 1997; Shishkina et al., 2018).
- 317 **Olivine-spinel oxygen barometer.** The oxygen fugacity of magmas was calculated 318 using the olivine-spinel oxygen barometer given by Ballhaus et al. (1991):

319
$$log_{10}fO_2(\Delta QFM) = 0.27 + 2505/T - 400P/T - 6log(X_{Fe}^{Ol}) - 3200(1 - X_{Fe}^{Ol})^2/T +$$

320
$$2\log(X_{Fe2+}^{Spl}) + 4\log(X_{Fe3+}^{Spl}) + 2630(X_{Al}^{Spl})^2/T$$
 (2)

where P is pressure in GPa, T is temperature in K, X_{Fe}^{Ol} is molar $Fe^{2+}/(Fe^{2+}+Mg^{2+})$ in 321 olivine, X_{Fe3+}^{Sp1} is molar Fe³⁺/ ΣR^{3+} in spinel, X_{A1}^{Sp1} is molar Al/ ΣR^{3+} in spinel, and X_{Fe2+}^{Sp1} is 322 molar $Fe^{2+}/(Fe^{2+}+Mg^{2+})$ in spinel. Olivine grains in the samples from the intrusions in the 323 CAOB have Fo contents varying from 82 to 90, with most being >84 (Table S1), and 324 325 those from the Dali picrite have Fo contents varying from 82 to 92 (Kamenetsky et al., 2012; Liu et al., 2017), which are all applicable to the equation. The pressure was 326 327 calculated using the clinopyroxene geobarometer given by Nimis and Ulmer (1998) (Table S1). The $Fe^{3+}/\Sigma Fe$ of the spinel from the Jinbulake, Erbutu, Baixintan, 328 329 Huangshannan and Tulaergen intrusions is corrected based on the EPMA data obtained in this study, whereas the $Fe^{3+}/\Sigma Fe$ of the spinel from the Heishan. Luodong intrusions and 330 Dali picrite cannot be corrected as the EPMA data were collected from the literature. The 331 magma fO_2 calculated using uncorrected Fe³⁺/ Σ Fe of the spinel is 0.2 to 0.6 log units 332 lower than that using corrected $Fe^{3+}/\Sigma Fe$ (Cao et al., 2019). However, the bias becomes 333 smaller with increasing fO_2 , which is <0.4 log units when fO_2 is >FMQ+1, and is <0.2 log 334 units when fO_2 is >FMQ+1.5 (Cao et al., 2019). 335

336 The accuracy of the results depends on whether or not the olivine-spinel pairs in the rocks are in chemical equilibrium (Ballhaus et al., 1991). The spinel grains in this study 337 overall are euhedral, fresh and homogeneous, and are commonly enclosed within olivine 338 339 (Fig. S1c). The textures showing chemical disequilibrium, such as complex zoning. embayment, symplectite and sieve texture, are not observed in both minerals. In addition, 340 341 the olivine-spinel pairs in the rocks from the intrusions in the CAOB overall have $ln Kd_{Mg/Fe}$ Ol-Spl positively correlated with XCr^{3+} [molar $Cr^{3+}/(Fe^{3+}+Cr^{3+}+Al^{3+})$] along the 342 equilibrium lines between 600 and 700°C (Fig. 7), indicating that the olivine-spinel pairs 343 344 reached chemical equilibrium. The temperatures of the equilibrium lines on Fig. 7 were estimated from the experimental data related to the reciprocal reaction (FeCr₂O₄ 345 $+MgAl_2O_4 = MgCr_2O_4 + FeAl_2O_4$ in spinel (Liermann and Ganguly, 2003), which are 346 347 consistent with the equilibrium temperatures calculated using the olivine-spinel thermometer given by Ballhaus et al. (1991) (Table S1). It is noted that the obtained 348 temperature values are the closure temperatures of Mg-Fe²⁺ diffusion between olivine 349 350 and spinel on subsolidus cooling, which are lower than the crystallization temperature of minerals (Kamenetsky et al., 2001). However, the fO_2 could be only elevated by ~0.2 log 351 units due to subsolidus Mg-F e^{2+} equilibrium between the olivine-spinel pairs (Birner et 352 al., 2018). Therefore, the fO_2 values obtained using the closure temperatures of the 353 354 olivine-spinel pairs can be taken as the magma fO_2 of the intrusions.

Using the equation 2, we obtained the magma fO_2 of the Jinbulake, Heishan, Erbutu, Baixintan, Huangshannan, Luodong and Tulaergen intrusions, which ranges from FMQ+1.2 to FMQ+2.6, FMQ+1.3 to FMQ+2.3, FMQ-0.1 to FMQ+1.2, FMQ+1.3 to FMQ+3.0, FMQ+0.6 to FMQ+2.6, FMQ+0.3 to FMQ+1.7, FMQ+2.5 to FMQ+2.9,

16

respectively (Table S1 and Fig. 8a). Although the values for the Heishan and Luodong intrusions were calculated using uncorrected $Fe^{3+}/\Sigma Fe$ of the spinel, the upper values should be reliable (*c.f.*, Cao et al., 2019). These data, together with the magma fO_2 of the Huangshandong, Huangshanxi, Poyi and Hongqiling No.1 and No.2 intrusions obtained in our earlier studies (Cao et al., 2019; Wei et al., 2019), display a negative correlation between the magma fO_2 and the Fo contents of olivine, except for the Erbutu intrusion (Fig. 8b).

The olivine-spinel pairs from the Dali picrite plot between the equilibrium lines at 900 and 1100°C (Fig. 7). The magma fO_2 of the Dali picrite varies from FMQ+0.2 to FMQ+0.8 (Fig. 8a). Given that the uncorrected Fe³⁺/ Σ Fe of the spinel was used in the calculation, the results could be underestimated by ~0.6 log units in this case (*c.f.*, Cao et al., 2019). However, even if the bias is considered, the magma fO_2 of the Dali picrite is still much lower than the magma fO_2 of the mafic-ultramafic intrusions in the CAOB (Fig. 8a).

Vanadium partitioning in olivine (D_V^{Ol}). Experimental results demonstrated that the partition coefficient of V between olivine and melt will decrease with elevating magma fO_2 (*e.g.*, Canil, 1997, 2002; Mallmann and O'Neill, 2013; Laubier et al., 2014; Shishkina et al., 2018). This relationship was used to calculate the magma fO_2 of hydrous arc basalts (Shishkina et al., 2018), *i.e.*,

378
$$\Delta FMQ = -3.07 \times log D_{v}^{Ol} - 3.34 \qquad (3)$$

A common way to measure D_{ν}^{Ol} is to acquire the V concentration of melt inclusion and host olivine in basalts. However, melt inclusions trapped in the olivine of cumulates 381 are difficult to be found and analyzed as they are usually very small. We therefore chose an alternative protocol to estimate the D_{ν}^{Ol} . 382

383 Vanadium and Sc are highly incompatible to olivine and have similar diffusion rates 384 between olivine and trapped liquid in crystal mush (Locmelis et al., 2019), the V/Sc ratio 385 of olivine is thus hardly affected by the trapped liquid shift effect. In addition, the V/Sc ratio of olivine is resistant to post-magmatic overprints, crustal contamination and 386 crystallization of small amounts of spinel (<5%) (Lee et al., 2005; Locmelis et al., 2019). 387 Nevertheless, we tried to analyze the core part of the best-preserved olivine grains in each 388 sample to warrant that the primary V/Sc ratio of olivine is acquired. In theory, the V/Sc 389 ratio of olivine can be calculated using the equation: 390

391
$$\left(\frac{V}{Sc}\right)_{Ol} = \frac{D_{\nu}^{Ol} \times V_{Liq}}{D_{Sc}^{Ol} \times Sc_{Liq}}$$
(4)

Since D_{Sc}^{Ol} is constant at ~0.2 (Villemant et al., 1981; Sun and Liang, 2013), the 392 equation 4 can be simplified as the equation: 393

394
$$\left(\frac{V}{Sc}\right)_{Ol} = \frac{D_{v}^{Ol} \times V_{Liq}}{0.2 \times Sc_{Liq}}$$
(5)

 D_V^{Ol} can be then acquired through the equation: 395

396
$$D_{\nu}^{Ol} = 0.2 \times \frac{\left(\frac{V}{Sc}\right)_{Ol}}{\left(\frac{V}{Sc}\right)_{Liq}} \quad (6)$$

397

If equation 6 is combined with equation 3, the magma fO_2 can be calculated by the 398 equation:

399
$$\Delta FMQ = -3.07 \times log \left[0.2 \times \frac{\left(\frac{V}{Sc}\right)_{Ol}}{\left(\frac{V}{Sc}\right)_{Liq}} \right] - 3.34 \qquad (7)$$

400 Although V and Sc are highly incompatible to both olivine and orthopyroxene, Sc is 401 more compatible to clinopyroxene than V (Canil, 2002). (V/Sc)_{Lig} would vary slightly 402 when olivine and/or orthopyroxene are on liquidus, but increase significantly when 403 clinopyroxene is on liquidus during the fractionation of mafic magmas (Laubier et al., 404 2014). Most samples in this study contain olivine and/or orthopyroxene as major cumulus 405 minerals (Fig. S1), except for those from the Jinbulake intrusion. Therefore, $(V/Sc)_{Lig}$ can 406 be referred to the V/Sc ratio of the primary magma for each intrusion in the CAOB 407 (Table S2), and then the magma fO_2 of the intrusions can be directly calculated using 408 equation 7 (Table S3).

Comparison of the results based on the two methods. The obtained magma fO_2 values based on the two methods are consistent with each other within uncertainties (Fig. 9a). The V/Sc ratios of the olivine from the Erbutu, Huangshannan, Hongqiling No.1 and No.2 intrusions generally decrease with increasing magma fO_2 values that were obtained based on the olivine-spinel oxygen barometer (Fig. 9b), indicating that the obtained magma fO_2 values in this study is reliable (*c.f.*, Canil, 1997, 2002; Mallmann and O'Neill, 2013; Laubier et al., 2014; Shishkina et al., 2018).

In summary, the magma fO_2 values of the arc-hosted Jinbulake and Heishan intrusions are comparable to those of the post-collisional Baixintan, Huangshandong, Huangshanxi, Huangshannan, Tulaergen, Hongqiling No.1 and No.2 intrusions. The magma fO_2 values of the mafic-ultramafic intrusions in the CAOB overall have a range similar to those of arc basalts (FMQ+0.5 to FMQ+6; Woodland et al., 2006), much higher than those of MORBs (FMQ-1 to FMQ+0.5; Cottrell and Kelley, 2011; Zhang et al., 2018) (Fig. 8a). The magma fO_2 values of the Erbutu, Poyi and Luodong intrusions

19

423 are lower than that of other intrusions in the CAOB, and overlap the upper fO_2 limit of 424 MORBs (Fig. 8a). In contrast, the magma fO_2 values of the Dali picrite are basically 425 within the range of MORBs (Fig. 8a).

426

427

442

443

DISCUSSIONS

428 The magma fO_2 of mafic-ultramatic intrusions in convergent margin settings could be 429 controlled by complex factors such as the oxidation and fertility states of the 430 metasomatized mantle sources (e.g., Rielli et al., 2017), and magmatic processes (e.g., 431 Lee et al., 2005). In this study, our results indicate that metasomatized mantle sources of 432 the mafic-ultramafic intrusions in the CAOB overall are slightly oxidized compared with 433 that of MORBs, and the elevated magma fO_2 of the intrusions in both arc and post-434 subduction, extensional settings is mainly attributed to the fractionation of hydrated 435 magmas derived from the metasomatized mantle.

436 Mantle fO₂ of the mafic-ultramafic intrusions in the CAOB

437 The arc-related Heishan intrusion and post-collisional Huangshannan, Poyi, Luodong 438 and Hongqiling No.2 intrusions have mantle fO_2 ranging from ~FMQ to ~FMQ+1.0 (Fig. 439 6), slightly higher than the mantle fO_2 (\leq FMQ) of MORBs (Frost and McCammon, 2008;

- Kelley and Cottrell, 2009, 2012; Rielli et al., 2018a), but much lower than the mantle fO_2
- 441 of arc basalts (FMQ+1 to FMQ+4, Woodland et al., 2006). These results indicate that the

mantle sources of mafic-ultramafic intrusions in the CAOB are not highly oxidized as

supposed for the subarc mantle. In addition, the mantle fO_2 is much lower than the

- 444 magma fO_2 of these intrusions (Fig. 8b), the high magma fO_2 of the intrusions in the
- 445 CAOB is thus not governed by the oxidation state of the mantle source alone.

446	The oxidation of the subarc mantle is attributed to the transportation of highly
447	oxidized, CO_3^{2-} , SO_4^{2-} , or Fe ³⁺ -rich fluids to the subarc mantle during subduction
448	(Mungall, 2002; Evans, 2006; Evans et al., 2012; Debret et al., 2016; Pons et al., 2016;
449	Debret and Sverjensky, 2017; Rielli et al., 2017). However, this process is dependent on
450	the subduction depth and temperature (Tomkins and Evans, 2015). Modeling results
451	indicate that sulfate tends to be released at shallower subduction zone and relatively low
452	temperatures, whereas sulfide tends to be released at deeper subduction zone and
453	relatively high temperatures (Tomkins and Evans, 2015). The mafic-ultramafic intrusions
454	in the CAOB are considered to have been derived from partial melts of the mantle wedge
455	in the spinel stability field (e.g., Zhang et al., 2016). It is likely that only minor slab-
456	derived, oxidized components was involved in the mantle wedge at this depth. In addition
457	the mantle sources of these intrusions in the CAOB are considered to have experienced
458	the interaction of the depleted lithospheric mantle with upwelling asthenospheric
459	materials due to slab break-off (Han et al., 2010; Li et al., 2012; Xie et al., 2012; Wei et
460	al., 2013; Mao et al., 2014, 2016; Deng et al., 2015). This process may also dilute the
461	oxidized components in the mantle wedge because asthenospheric materials are typically
462	more reduced than the lithospheric mantle by $\sim 1 \log$ unit (Wood et al., 1990). Therefore,
463	the mafic-ultramafic intrusions in the CAOB overall have mantle fO_2 slightly higher than
464	that for the mantle of MORBs.

465 Fractionation of hydrated magmas derived from metasomatized mantle sources

Experimental results indicate that the fractionation of olivine and clinopyroxene may slightly increase the $Fe^{3+}/\Sigma Fe$ of magmas and have a limited effect on the oxidization states of magmas (Cottrell and Kelley, 2011; Kelley and Cottrell, 2012). However, water

in silicate magmas can play an efficient 'catalyst' to promote the oxidation states of magmas if it is partially dissociated and loss H^+ at high temperatures (Carmichael, 1991; Cornejo and Mahood, 1997), or exsolved from the melt that carried more Fe^{2+} than Fe^{3+} (Bell and Simon, 2011). Mafic magmas tend to become more hydrous with fractionation because volatiles (*e.g.*, H₂O) are essentially incompatible to olivine and clinopyroxene. Therefore, the fractionation process could significantly elevate the oxidation states of hydrated, mafic magmas.

The mafic-ultramafic intrusions in the CAOB contain abundant hydrous minerals 476 477 such as amphibole and phlogopite (e.g., Deng et al., 2014; Su et al., 2011; Xie et al., 2012; Wei et al., 2013, 2015). On the plot of Alz versus TiO₂, the clinopyroxene from the 478 479 intrusions in the CAOB has Alz/Ti scattered along the arc cumulate trend, in contrast to 480 the low Alz/Ti of the clinopyroxene from the sulfide-bearing mafic-ultramafic intrusions 481 in the Emeishan LIP (Fig. 10). The high Alz values of the clinopyroxene from the CAOB 482 are attributed to the idea that more Al would enter the tetrahedral site of clinopyroxene with increasing H_2O content of melt (c.f., Loucks, 1990). This is consistent with an 483 484 interpretation that the parental magmas of the intrusions in the CAOB may be hydrated 485 due to the derivation from the mantle sources metasomatized by slab-derived melts/fluids. 486 There is an overall negative correlation between the magma fO_2 and the Fo contents of 487 olivine for the intrusions in the CAOB (Fig. 8b), showing that the magmas became more 488 oxidized with fractionation. Therefore, the H₂O content of magmas derived from the 489 metasomatized mantle and relative degrees of the fractionation of magmas are likely two 490 key factors controlling magma fO_2 of the mafic-ultramafic intrusions in convergent 491 margin settings.

492 The Erbutu intrusion is an exceptive case as the olivine grains of the intrusion have 493 Fo contents comparable with those for the olivine of the Jinbulake and Heishan intrusions, but the intrusion has much lower magma fO_2 than the latter two intrusions (Fig. 8b). The 494 495 parental magma of the Erbutu intrusion is thought to be boninitic that may have been 496 emplaced early in the subduction history (c.f., Jian et al., 2010; Peng et al., 2013). As the 497 oxidation of the mantle wedge by the metasomatizing agents could occur after subduction 498 initiation in 1 Myr. (c.f., Brounce et al., 2015), it is likely that the mantle source of the 499 Erbutu intrusion is relatively reduced, thus the magma fO_2 of this intrusion is lower than 500 that of other intrusions in the CAOB for a given degree of fractionation of magma.

501

502 Magma fO₂ constraints for Ni-Cu sulfide mineralization in convergent margin 503 settings

Experimental results show that the sulfur solubility of silicate magma could increase 504 505 by an order of magnitude if the magma fO_2 increases from FMQ+0.5 to FMQ+1.5 (Luhr, 506 1990; Jugo et al., 2005; Jugo, 2009; Jugo et al., 2010). The mantle-derived mafic magmas 507 in intraplate settings usually have magma fO_2 ranging from FMQ-1 to FMQ+0.5 and 508 could dissolve a maximum of ~ 1500 ppm S (*c.f.*, Wood et al., 1990; Jugo et al., 2010), 509 therefore the formation of economic Ni-Cu sulfide deposits often requires the addition of 510 external crustal sulfur into the magmas (e.g., Li et al., 2001; Ripley and Li, 2003; Barnes 511 and Lightfoot, 2005; Wang et al., 2006; Mungall and Naldrett, 2008; Keays and Lightfoot, 512 2010; Taranovic et al., 2018). For instance, the Ni-Cu sulfide deposits in the Emeishan 513 LIP and the Jinchuan Ni-Cu deposit formed in a rifting setting have magma fO_2 overlapping with the range of MORBs, and the sulfides from the deposits have highly 514

variable δ^{34} S (-4 to +8‰, Fig. 11), indicating substantial addition of external crustal sulfur in the formation of these deposits (Ripley et al., 2005; Duan et al., 2016; Wang et al., 2018).

In contrast, the mantle-derived mafic magmas in convergent margin settings have fO_2 518 ranging from FMQ+0.5 to FMQ+3 (Fig. 8a) and could dissolve ~1800 to ~13,000 ppm S 519 520 (Jugo et al., 2010), much higher than the S solubility of the magmas in intraplate settings. In addition, the sulfides from the Ni-Cu sulfide-bearing mafic-ultramafic intrusions in the 521 post-subduction, extensional setting in the CAOB have δ^{34} S values (-1.0 to +1.3‰) 522 nearly identical to that of the MORB mantle (Fig. 11), despite the large δ^{34} S range (-10.0 523 to +5.4%) of the sulfides from the metasomatized mantle xenoliths (Rielli et al., 2018b). 524 525 This was interpreted as the magmas of the Ni-Cu sulfide-bearing mafic-ultramafic 526 intrusions in the CAOB contain dominantly mantle-derived sulfur with trivial addition of external crustal sulfur (Wei et al., 2019). Therefore, the high magma fO_2 and the MORB 527 mantle-like δ^{34} S of the mafic-ultramatic intrusion in the CAOB indicate that highly 528 oxidized, mantle-derived magmas may be capable of dissolving enough mantle-derived 529 530 sulfur to form magmatic Ni-Cu sulfide deposits so that the addition of external crustal 531 sulfur is not always necessary in such cases. In addition, the mafic-ultramafic intrusions in the CAOB that have sulfides with mantle-like δ^{34} S values generally have magma 532 fO_2 >FMQ+1, whereas the Erbutu intrusion that has sulfides with the highest $\delta^{34}S$ values 533 534 has magma $fO_2 \leq FMQ+1$ (Fig. 11), we thus consider that the mantle-derived matic magmas with fO_2 greater than ~FMQ+1.0 may be able to dissolve sufficient mantle-535 derived sulfur to form important Ni-Cu sulfide deposits in convergent margin settings 536 537 (*c.f.*, Rielli et al., 2018a).

538 On the other hand, the formation of economic Ni-Cu sulfide deposits from the highly oxidized, mantle-derived magmas depends on how the magmas can be reduced to reach 539 540 sulfide saturation so that the sulfide melts can be segregated from the magmas (Tomkins 541 et al., 2012). This can be examined by comparing the fO_2 between the parental magmas prior to sulfide saturation and the magmas concurrent with sulfide saturation (e.g., Wei et 542 543 al., 2019). The magma fO_2 obtained by the olivine-spinel oxygen barometer in this study 544 can represent the parental magma fO_2 before sulfide saturation. The fO_2 of the magmas 545 concurrent with sulfide saturation for the intrusions in the CAOB were estimated using 546 Fe-Ni exchange between olivine and sulfide liquid (e.g., Feng et al., 2017; Mao et al., 2018; Wei et al., 2019). As shown in Fig. 12, the magma fO_2 at sulfide saturation is 547 considerably lower than the fO_2 of parental magmas for each intrusion, indicating that the 548 549 oxidized magmas was indeed reduced with the sulfide saturation of magmas. A possible way to trigger the reduction is the crystallization of magnetite (Jenner et al., 2010). 550 551 However, this mechanism does not appear as the driver of magma reduction in the CAOB 552 because the examined rocks in this study contain few magnetite. Alternatively, the reduction of oxidized magmas can be triggered by the addition of organic-carbon or 553 554 graphite-rich sedimentary rocks, which was evidenced by the C isotope studies on a few intrusions in the CAOB (e.g., Wei et al., 2019) and the O isotope studies of the olivine in 555 the lower zone of the Huangshanxi intrusion (Mao et al., 2019). 556

- 557
- 558

IMPLICATIONS

559 Most Ni-Cu sulfide-bearing mafic-ultramafic intrusions in the CAOB have magma 560 fO_2 (FMQ+0.5 to FMQ+3) much higher than that of MORBs (FMQ-1 to FMQ+0.5),

561 consistent with the global observation that the mafic-ultramafic intrusions emplaced in convergent margin settings have relatively high magma fO_2 . In contrast, the mantle fO_2 of 562 563 these intrusions ranges from FMQ to \sim FMQ+1.0, just slightly higher than that of MORBs 564 $(\leq$ FMQ). Because the amounts of oxidized components that were added to the metasomatized mantle wedges generally decrease with the depth of the mantle wedges in 565 convergent margin settings, the slightly oxidized mantle source of the intrusions in the 566 567 CAOB is likely related to the limited amounts of slab-derived, oxidized components added to mantle wedges and relatively deep mantle wedges where the partial melting 568 569 occurred. The negative correlation of the magma fO_2 and the Fo contents of the olivine of the intrusions in the CAOB indicates that the magma fO_2 could be elevated with the 570 571 fractionation of hydrated, mafic magmas derived from metasomatized mantle sources. In 572 addition, the mafic-ultramafic intrusions that host economic Ni-Cu sulfide deposits in the CAOB usually have sulfides with mantle-like δ^{34} S (-1.0 to +1.1‰) and magma 573 574 $fO_2 > FMQ+1$, indicating that the relatively oxidized magmas may be capable of 575 dissolving enough mantle-derived sulfur to form economic Ni-Cu sulfide deposits in convergent margin settings. The sulfide saturation of the oxidized, mafic magmas may be 576 577 triggered by the addition of organic-carbon or graphite-rich sedimentary rocks into the magmas. Therefore, our results imply that the addition of external crustal sulfur is not so 578 579 compulsory to trigger the sulfide saturation of highly oxidized, mantle-derived mafic 580 magmas and the formation of economic Ni-Cu sulfide deposits in convergent margin 581 settings, although it is very important in the formation of giant Ni-Cu sulfide deposits 582 such as those at Noril'sk in Russia (Ripley and Li, 2013).

583

~	01
2	XД
2	UT

ACKNOWLEDGEMENTS AND FUNDING

585	This work was supported by grants from the National Natural Science Foundation of
586	China (No. 41730423 and 41902077), and China Postdoctoral Science Foundation Grant
587	(No. 2019M653103). Shenghong Yang provided the samples of the Jinbulake intrusion
588	and Benxun Su shared the EPMA data for the olivine and spinel of the Luodong intrusion.
589	Constructive reviews by Andrew Tomkins and an anonymous reviewer have greatly
590	improved the quality of this manuscript.
591	
592	REFERENCES
593	Ballhaus C., Berry R.F., and Green D.H. (1991) High pressure experimental calibration
594	of the olivine-orthopyroxene-spinel oxygen geobarometer: implications for the
595	oxidation state of the upper mantle. Contributions to Mineralogy and Petrology 107,
596	27–40.
597	Barnes SJ., and Lightfoot P.C. (2005) Formation of magmatic nickel-sulfide ore
598	deposits and processes affecting their copper and platinum-group element contents.
599	In Economic Geology 100th Anniversary Volume (eds. J. W. Hedenquist, J. H.
600	Thompson, R. J. Goldfarb and J. P. Richards). pp. 179-213.
601	Bell A.S., and Simon A. (2011) Experimental evidence for the alteration of the Fe ³⁺ / Σ Fe
602	of silicate melt caused by the degassing of chlorine-bearing aqueous volatiles.
603	Geology 39, 499–502.
604	Bénard A., Woodland A.B., Arculus R.J., Nebel O., and McAlpine S.R.B. (2018)
605	Variation in sub-arc mantle oxygen fugacity during partial melting recorded in
606	refractory peridotite xenoliths from the West Bismarck Arc. Chemical Geology 486,
607	16–30.
608	Birner A., Cottrell E., Warren J.M., Kelley K.A., Davis F.A. (2018) Peridotites and
609	basalts reveal broad congruence between two independent records of mantle fO_2
610	despite local redox heterogeneity. Earth and Planetary Science Letters 494, 172-189.
611	Bottinga Y., and Allegre C. (1976). Geophysical, petrological and geochemical models of
612	the oceanic lithosphere. Tectonophysics 32, 9-59.

613 Brounce M., Kellev K.A., Cottrell E. and Reagan M.K. (2015) Temporal evolution of mantle wedge oxygen fugacity during subduction initiation. Geology 43, 775–778. 614 615 Brounce M., Stolper E., and Eiler J. (2017) Redox variations in Mauna Kea lavas, the oxygen fugacity of the Hawaiian plume, and the role of volcanic gases in Earth's 616 oxygenation. PNAS 114, 8997-9002. 617 Canil D. (1997) Vanadium partitioning and the oxidation state of Archean komatiite 618 619 magmas. Nature 389, 842-845. Canil D. (2002) Vanadium in peridotites, mantle redox and tectonic environments: 620 621 Archean to present. Earth and Planetary Science Letters 195, 75-90. 622 Canil D., and Fedortchouk Y. (2001) Olivine-liquid partitioning of vanadium and other 623 trace elements, with applications to modern and ancient picrites. The Canadian Mineralogist 39, 319-330. 624 Cao Y., Wang C.Y., and Wei B. (2019) Magma oxygen fugacity of Permian to Triassic 625 626 Ni-Cu sulfide-bearing mafic-ultramafic intrusions in the central Asian orogenic belt, North China. Journal of Asian Earth Sciences 173, 250-262. 627 628 Carmichael I.S.E. (1991) The redox states of basic and silicic magmas- a reflection of their source regions. Contributions to Mineralogy and Petrology 106, 129–141. 629 Cornejo P.C., and Mahood G.A. (1997) Seeing past the effects of re-equilibration to 630 631 reconstruct magmatic gradients in plutons: La Gloria Pluton, central Chilean Andes. 632 Contributions to Mineralogy and Petrology 127, 159–175. Cottrell E., and Kelley K.A. (2011) The oxidation state of Fe in MORB glasses and the 633 634 oxygen fugacity of the upper mantle. Earth and Planetary Science Letters 305, 270– 282. 635 636 Dauphas N., Teng F.-Z., and Arndt N.T. (2010) Magnesium and iron isotopes in 2.7 Ga Alexo komatiites: Mantle signatures, no evidence for Soret diffusion, and 637 638 identification of diffusive transport in zoned olivine. Geochimica et Cosmochimica 639 Acta 74, 3274–3291. 640 Debret B., and Sverjensky D.A. (2017) Highly oxidising fluids generated during serpentinite breakdown in subduction zones. Scientific Report 1-6. 641 Debret B., Millet M.-A., Pons M.-L., Bouilhol P., Inglis E., and Williams H. (2016) 642 Isotopic evidence for iron mobility during subduction. Geology 44, 215-218. 643

- 644 Deng Y.F., Song X.Y., Chen L., Zhou T., Pirajno F., Yuan F., Xie W., and Zhang D.
- 645 (2014) Geochemistry of the Huangshandong Ni-Cu deposit in northwestern China:

Implications for the formation of magmatic sulfide mineralization in orogenic belts.
Ore Geology Reviews 56, 181–198.

- Deng Y.F., Song X.Y., Hollings P., Zhou T.F., Yuan F., and Zhang D. (2015) Role of
 asthenosphere and lithosphere in the genesis of the Early Permian Huangshan maficultramafic intrusion in the Northern Tianshan, NW China. Lithos 227, 241–254.
- Duan J., Li C., Qian Z., Jiao J., Ripley E.M., and Feng Y. (2016) Multiple S isotopes,
 zircon Hf isotopes, whole-rock Sr-Nd isotopes, and spatial variations of PGE tenors
- 653 in the Jinchuan Ni-Cu-PGE deposit, NW China. Mineralium Deposita 51, 557–574.
- Evans B.W., Dyar M.D., and Kuehner S.M. (2012) Implications of ferrous and ferric iron
 in antigorite. American Mineralogist 97, 184–196.
- Evans K.A. (2006) Redox decoupling and redox budgets: Conceptual tools for the study
 of earth systems. Geology 34, 489-492.
- Feng Y., Qian Z., Xu G., Duan J., Chen B., Sun T., Jiang C., and Ren M. (2017) Rockforming mineral features of Permian mineralized mafic-ultramafic intrusions in East
 Tianshan Mountains and their implications for intrusion generation. Acta
 Petrological et Mineralogical 36, 519-534 (in Chinese with English abstract).
- Frost D.J., and McCammon C.A. (2008) The redox state of earth's mantle. Annual
 Review of Earth and Planetary Science 389-420.
- Gaillard F., Scaillet B., Pichavant M., and Lacono-Marziano G. (2015) The redox
 geodynamics linking basalts and their mantle sources through space and time.
 Chemical Geology 418, 217–233.
- Gao J.-F., Zhou M.-F., Lightfoot P., and Qu W. (2012) Heterogeneous Os isotope
 compositions in the Kalatongke sulfide deposit, NW China: the role of crustal
 contamination. Mineralium Deposita 47, 731–738.
- Godel B., Barnes S.J., and Maier W. D. (2011) Parental magma composition inferred
 from trace element in cumulus and intercumulus silicate minerals: an example from
 the lower and lower critical zones of the Bushveld Complex, South-Africa. Lithos
 125, 537–552.

- Han C., Xiao W., Zhao G., Ao S., Zhang J., Qu W., and Du A. (2010) In-situ U-Pb, Hf
- and Re-Os isotopic analyses of the Xiangshan Ni-Cu- Co deposit in Eastern Tianshan (Xinjiang), Central Asia Orogenic Belt: Constraints on the timing and
- 677 genesis of the mineralization. Lithos 120, 547–562.
- Han C., Xiao W., Zhao G., Qu W., and Du A. (2007) Re–Os dating of the Kalatongke
 Cu–Ni deposit, Altay Shan, NW China, and resulting geodynamic implications. Ore
 Geology Reviews 32, 452–468.
- Hao L., Zhao X., Boorder H.D., Lu J., Zhao, Y., and Wei, Q. (2014) Origin of PGE
 depletion of Triassic magmatic Cu-Ni sulfide deposits in the central-southern area of
 Jilin province, NE China. Ore Geology Reviews 63, 226–237.
- Ionov D.A., and Wood B.J. (1992) The oxidation state of subcontinental mantle: oxygen
 thermobarometry of mantle xenoliths from central Asia. Contributions to
 Mineralogy and Petrology 111, 179–193.
- Jahn B.M., Windley B., Natal'in B., and Dobretsov N. (2004) Phanerozoic continental
 growth in Central Asia. Journal of Asian Earth Sciences 23, 599-603.
- Jahn B.M., Wu F.Y., and Chen B. (2000) Massive granitoid generation in Central Asia:
- 690 Nd isotope evidence and implication for continental growth in the Phanerozoic.
 691 Episodes 23, 82–92.
- Jenner F.E., O'Neill H.St.C., Arculus R.J., and Mavrogenes J.A. (2010) The magnetite
 crisis in the evolution of arc-related magmas and the initial concentration of Au, Ag
 and Cu. Journal of Petrology 51, 2445–2464.
- Jian P., Liu D., Kröner A., Windley B.F., Shi Y., Zhang W., Zhang F., Miao L., Zhang L.,
 and Tomurhuu D. (2010) Evolution of a Permian intraoceanic arc-trench system in
 the Solonker suture zone, Central Asian Orogenic Belt, China and Mongolia. Lithos
 118, 169-190.
- Jiang Y.-H., Jiang S.-Y., Dai B.-Z., Liao S.-Y., Zhao K.-D., and Ling H.-F. (2009)
 Middle to late Jurassic felsic and mafic magmatism in southern Hunan Province,
 southeast China: Implications for a continental arc to rifting. Lithos 107, 185–204.
- Jugo P.J. (2009) Sulfur content at sulfide saturation in oxidized magmas. Geology 37,
 415–418.

Jugo P.J., Luth R.W., and Richards J.P. (2005) Experimental data on the speciation of					
sulfur as a function of oxygen fugacity in basaltic melts. Geochimica et					
Cosmochimica Acta 69, 497–503.					
Jugo P.J., Wilke M., and Botcharnikov R.E. (2010) Sulfur K-edge XANES analysis of					
natural and synthetic basaltic glasses: Implications for S speciation and S content as					
function of oxygen fugacity. Geochimica et Cosmochimica Acta 74, 5926-5938.					
Kamenetsky V.S., Chung SL., Kamenetsky M.B., and Kuzmin D.V. (2012) Picrites					
from the Emeishan large igneous province, SW China: a compositional continuum					
in primitive magmas and their respective mantle sources. Journal of Petrology 53,					
2095-2113.					
Kamenetsky V.S., Crawford A.J., and Meffre S. (2001) Factors controlling chemistry of					
magmatic spinel: An empirical study of associated olivine, Cr-spinel and melt in-					
clusions from primitive rocks. Journal of Petrology 42, 655-671.					
Keays R.R., and Lightfoot P.C. (2010) Crustal sulfur is required to form magmatic Ni-Cu					
sulphide deposits; evidence from chalcophile element signatures of Siberian and					
Deccan Trap basalts. Mineralium Deposita 45, 241–257					
Kelley K.A., and Cottrell E. (2009) Water and the oxidation state of Subduction Zone					
Magmas. Science 325, 605–607.					
Kelley K.A., and Cottrell E. (2012) The influence of magmatic differentiation on the					
oxidation state of Fe in a basaltic arc magma. Earth and Planetary Science Letters					
329–330, 109–121.					
Kelley K.A., Plank T., Grove T.L., Stolper E.M., Newman S., and Hauri E. (2006).					
Mantle melting as a function of water content beneath back-arc basins. Journal of					
Geophysical Research: Solid Earth 111, B09208.					
Kelley K.A., Plank T., Newman S., Stolper E., Grove T.L., Parman S., and Hauri E.					
(2010) Mantle melting as a function of water content beneath the Mariana arc.					
Journal of Petrology 51, 1711–1738.					
Kress V.C., and Carmichael I.S.E. (1991) The compressibility of silicate liquids					
containing Fe ₂ O ₃ and the effect of composition, temperature, oxygen fugacity and					
pressure on their redox states. Contributions to Mineralogy and Petrology108, 82-92					

734 Labidi J., Cartigny P., and Moreira M. (2013) Non-chondritic sulphur isotope composition of the terrestrial mantle. Nature 501, 208–211. 735 736 Labidi J., Cartigny P., Hamelin C., Moreira M., and Dosso L. (2014) Sulfur isotope budget (³²S, ³³S, ³⁴S and ³⁶S) in Pacific-Antarctic Ridge basalts; a record of mantle 737 source heterogeneity and hydrothermal sulfide assimilation. Geochimica et 738 739 Cosmochimica Acta133, 47-67. 740 Laubier M., Grove T.L., and Langmuir C.H. (2014) Trace element mineral/melt 741 partitioning for basaltic and basaltic andesitic melts: An experimental and laser ICP-742 MS study with application to the oxidation state of mantle source regions. Earth and 743 Planetary Science Letters 392, 265–278. Lee C.T.A., Leeman W.P., Canil D., and Li Z.X.A. (2005) Similar V/Sc systematics in 744 MORB and arc basalts: Implications for the oxygen fugacities of their mantle source 745 regions. Journal of Petrology 46, 2313-2336. 746 Lee C.T.A., Luffi P., Le Roux V., Dasgupta R., Albaréde F., and Leeman W.P. (2010) 747 The redox state of arc mantle using Zn/Fe systematics. Nature 468, 681–685. 748 749 Li C., and Ripley E.M. (2011) The giant Jinchuan Ni-Cu-(PGE) deposit; tectonic setting, magma evolution, ore genesis, and exploration implications. Reviews in Economic 750 751 Geology 17, 163–180. Li C., Maier W.D., and Waal S.A. (2001) Magmatic Ni-Cu versus PGE deposits: 752 753 contrasting genetic controls and exploration implication. South African Journal of 754 Geology 104, 205–214. 755 Li C., Zhang M.J., Fu P., Qian Z.Z., Hu P.Q., and Ripley E.M. (2012) The Kalatongke magmatic Ni-Cu deposits in the Central Asian Orogenic Belt, NW China: product of 756 757 slab window magmatism? Mineralium Deposita 47, 51-67. Li C., Zhang Z., Li W., Wang Y., Sun T., and Ripley E.M. (2015) Geochronology, 758 759 petrology and Hf-S isotope geochemistry of the newly-discovered Xiarihamu 760 magmatic Ni-Cu sulfide deposit in the Qinghai-Tibet plateau, western China. Lithos 761 216, 224-240. Li J.L., Schwarzenbach E.M., John T., Ague J.J., Huang F., Gao, J., Klemd R., 762 763 Whitehouse M.J., and Wang X.S. (2020) Uncovering and quantifying the subduction 764 zone sulfur cycle from the slab perspective. Nature Communications 11, 1-12.

Li Y. (2018) Temperature and pressure effects on the partitioning of V and Sc between
 clinopyroxene and silicate melt: Implications for mantle oxygen fugacity. American
 Mineralogist 103, 819–823.

- Liermann H.P., and Ganguly J. (2003) Fe²⁺-Mg fractionation between orthopyroxene and
 spinel: experimental calibration in the system FeO–MgO–Al₂O₃–Cr₂O₃–SiO₂, and
 applications. Contributions to Mineralogy and Petrology 154, 217–227.
- Liu J., Xia Q.-K., Kuritani T., Hanski E., and Yu H.-R. (2017) Mantle hydration and the
 role of water in the generation of large igneous provinces. Nature Communication 8,
 1824.
- Locmelis M., Arevalo R.D., Puchtel I.S., Fiorentini M.L., and Nisbet E.G. (2019)
 Transition metals in komatiitic olivine: Proxies for mantle composition, redox
 conditions, and sulfide mineralization potential. American Mineralogist 104, 1143–
 1155.
- Loucks R.R. (1990) Discrimination from opholitic and nonopholitic ultramafic–mafic
 allochthons in orogenic belts by the Al/Ti ratios in clinopyroxene. Geology 18, 346–
 349.
- Luhr J.F. (1990) Experimental phase relations of water- and sulfur-saturated arc magmas
 and the 1982 eruptions of El Chichon volcano. Journal of Petrology 31, 1071–1114.
- Maier W.D., Barnes S.J., Chinyepi G., Barton J.J., Eglington B., and Setshedi T. (2008)
 The composition of magmatic Ni-Cu-(PGE) sulfide deposits in the Tati and SelebiPhikwe belts of eastern Botswana. Mineralium Deposita 43, 37-60.
- Mallmann G., and O'Neill H.S.C. (2009) The crystal/melt partitioning of V during
 mantle melting as a function of oxygen fugacity compared with some other elements
 (Al, D, Ca, Sa, Ti, Ca, Va, Za and Nb) Jacomel of Dataslass 50, 17(5, 1704)
- (Al, P, Ca, Sc, Ti, Cr, Fe, Ga, Y, Zr and Nb). Journal of Petrology 50, 1765–1794.
- Mallmann G., and O'Neill H.S.C. (2013) Calibration of an empirical thermometer and
 oxybarometer based on the partitioning of Sc, Y and V between olivine and silicate
 melt. Journal of Petrology 54, 933–949.
- Manor M.J., Scoates J.S., Nixon G.T., and Ames D.E. (2016) The giant mascot Ni-Cu PGE deposit, British Columbia: mineralized conduits in a convergent margin
 tectonic setting. Economic Geology 111, 57–87.

- Mao Y., Qin K., Barnes S. J., Ferraina C., Iacono–Marziano G., Verrall M., Tang D., and
 Xue S. (2018) A revised oxygen barometry in sulfide-saturated magmas and
 application to the Permian magmatic Ni–Cu deposits in the southern Central Asian
 Orogenic Belt. Mineralium Deposita 53, 731-755.
- Mao Y., Qin K., Li C., and Tang D. (2015) A modified genetic model for the
 Huangshandong magmatic sulfide deposit in the Central Asian Orogenic Belt,
 Xinjiang, western China. Mineralium Deposita 50, 65–82.
- Mao Y., Qin K., Li C., Xue S., and Ripley E.M. (2014) Petrogenesis and ore genesis of
 the Permian Huangshanxi sulfide ore-bearing mafic-ultramafic intrusion in the
 Central Asian Orogenic Belt, western China. Lithos 200–201, 111–125.
- Mao Y., Qin K., Tang D., Feng H. and Xue S. (2016) Crustal contamination and sulfide
 immiscibility history of the Permian Huangshannan magmatic Ni-Cu sulfide deposit,
 East Tianshan, NW China. Journal of Asian Earth Sciences 129, 22–37.
- Mao Y.J., Barnes S.J., Qin K.Z., Tang D., Martin L., Su B., and Evans N.J. (2019) Rapid
 orthopyroxene growth induced by silica assimilation: constraints from sector-zoned
 orthopyroxene, olivine oxygen isotopes and trace element variations in the
 Huangshanxi Ni–Cu deposit, Northwest China. Contributions to Mineralogy and
 Petrology 174, 1–24.
- Mungall J. E. (2002) Roasting the mantle: Slab melting and the genesis of major Au and
 Au-rich Cu deposits. Geology 30, 915-918.
- Mungall J.E., and Naldrett A.J. (2008) Ore Deposits of the Platinum-Group Elements.
 Element 4, 253–258.
- Naldrett A.J. (2004) Magmatic sulfide deposits: geology, geochemistry and exploration.
 Springer, New York.
- Nilsson K., and Peach C.L. (1993) Sulfur speciation, oxidation state and sulfur
 concentration in backarc magmas. Geochimica et Cosmochimica Acta 57, 3807–
 3813.
- Nimis P., and Ulmer P. (1998) Clinopyroxene geobarometry of magmatic rocks Part 1:
 An expanded structural geobarometer for anhydrous and hydrous, basic and
 ultrabasic systems. Contributions to Mineralogy and Petrology 133, 122–135.

- Peng R., Zhai Y., Li C., and Ripley M. (2013) The Erbutu Ni-Cu deposit in the Central
 Asian Orogenic Belt: A Permian Magmatic Sulfide deposit related to boninitic
 magmatism in an arc setting. Economic Geology 108, 1879-1888.
- Pons M.-L., Debret B., Bouilhol P., Delacour A., and Williams H. (2016) Zinc isotope
 evidence for sulfate-rich fluid transfer across subduction zones. Nature
 Communication 7, 13794.
- Qin K.Z., Su B.X., Li X.H., Tang D.M., Sakyi P.A., Sun H., Xiao Q.H., and Liu P.P.
 (2011). SIMS zircon U-Pb geochronology and Sr-Nd isotopes of mafic-ultramafic
 intrusions in eastern Tianshan and Beishan in correlation with flood basalts in Tarim
 basin (NW China): Constraints on a 280 Ma mantle plume. American Journal of
 Sciences 29, 275–289.
- Rielli A., Tomkins A.G., Nebel O., Brugger J., Etschmann B., and Paterson D. (2018a)
 Garnet peridotites reveal spatial and temporal changes in the oxidation potential of
 subduction. Scientific Reports 8, 16411.
- Rielli A., Tomkins A.G., Nebel O., Brugger J., Etschmann B., Zhong R., Yaxley G.M.,
 and Paterson D. (2017) Evidence of sub-arc mantle oxidation by sulphur and carbon.
 Geochemical Perspective Letters 124–132.
- Rielli A., Tomkins A.G., Nebel O., Raveggi M., Jeon H., Martin L., and Ávila J.N.
 (2018b) Sulfur isotope and PGE systematics of metasomatised mantle wedge. Earth
 and Planetary Science Letters 497, 181–192.
- Ripley E.M., and Li C. (2003) Sulfur isotope exchange and metal enrichment in the
 formation of magmatic Cu-Ni-(PGE) deposits. Economic Geology 99, 635–641.
- Ripley E.M., and Li, C. (2013) Sulfide saturation in mafic magmas: Is external sulfur
 required for magmatic Ni-Cu-(PGE) ore genesis? Economic Geology 108, 45-58.
- Ripley E.M., Sarkar A., and Li C. (2005) Mineralogical and stable isotope studies of
 hydrothermal alteration at the Jinchuan Ni-Cu deposit. Economic Geology 100,
 1349–1361.
- Sengör A.C.A., Natal'in B.A., and Burtmann V.S. (1993) Evolution of the Altaid tectonic
 collage and Palaeozoic crustal growth in Eurasia. Nature 364, 299–306.
- 854 Shishkina T.A., Portnyagin M.V., Botcharnikov R.E., Almeev R.R., Simonyan A.V.,
- 655 Garbe-Schönberg D., Schuth S., Oeser M., and Holtz F. (2018) Experimental

856 Calibration and Implications of Olivine-Melt Vanadium Oxybarometry for Hydrous Basaltic Arc Magmas. American Mineralogist 103, 369-383. 857 858 Song X.-Y., and Li X.-R. (2009) Geochemistry of the Kalatongke Ni-Cu- (PGE) sulfide deposit, NW China: Implications for the formation of magmatic sulfide 859 mineralization in a postcollisional environment. Mineralium Deposita 44, 303–327. 860 Song X.-Y., Chen L.-M., Deng Y.-F., and Xie W. (2013) Syncollisional tholeiitic 861 862 magmatism induced by asthenosphere upwelling owing to slab detachment at the 863 southern margin of the Central Asian orogenic belt. Journal of Geological Society 864 170, 941-950. Song X.-Y., Yi J., Chen L., She Y., Liu C., Dang X.F., Yang Q., and Wu S. (2016) The 865 Giant Xiarihamu Ni-Co sulfide deposit in the East Kunlun Orogenic Belt, Northern 866 Tibet Plateau. China. Economic Geology 111, 29–55. 867 Su B.X., Qin K.Z., Lu Y., Sun H., and Sakyi P.A. (2015) Decoupling of whole-rock Nd-868 Hf and zircon Hf-O isotopic compositions of a 284 Ma mafic-ultramafic intrusion in 869 the Beishan Terrane, NW China. International Journal of Earth Sciences 104, 1721-870 871 1737. Su B.X., Qin K.Z., Sakyi P.A., Li X.H., Yang Y.H., Sun H., Tang D.M., Liu P.P., Xiao 872 Q.H., and Malaviarachchi S.P.K. (2011) U-Pb ages and Hf-O isotopes of zircons 873 874 from Late Paleozoic mafic-ultramafic units in the southern Central Asian Orogenic 875 Belt: tectonic implications and evidence for an Early-Permian mantle plume. Gondwana Research 20, 516–531. 876 Sun C., and Liang Y. (2013) The importance of crystal chemistry on REE partitioning 877 878 between mantle minerals (garnet, clinopyroxene, orthopyroxene, and olivine) and 879 basaltic melts. Chemical Geology 358, 23-36. Sun S.S., and McDonough W.F. (1989) Chemical and isotopic systematics of oceanic 880 881 basalts: implications for mantle composition and processes. Geological Society, 882 London, Special Publications 42, 313–345. 883 Sun T., Qian Z.-Z., Li C., Xia M.-Z., and Yang S.-H. (2013) Petrogenesis and economic potential of the Erhongwa mafic-ultramafic intrusion in the Central Asian Orogenic 884 Belt, NW China: Constraints from olivine chemistry, U-Pb age and Hf isotopes of 885 zircons, and whole-rock Sr-Nd-Pb isotopes. Lithos 182-183, 185-199. 886

887	Tang D., Qin K., Sun H., Su B., and Xiao Q. (2012) The role of crustal contamination in
888	the formation of Ni-Cu sulfide deposits in Eastern Tianshan, Xinjiang, Northwest
889	China: Evidence from trace element geochemistry, Re-Os, Sr-Nd, zircon Hf- O, and
890	sulfur isotopes. Journal of Asian Earth Sciences 49, 145-160.
891	Taranovic V., Ripley E. M., Li C. and Shirey, S.B. (2018) S, O, and Re-Os isotope
892	studies of the Tamarack Igneous Complex: melt-rock interaction during the early
893	stage of Midcontinent Rift Development. Economic Geology 113, 1161-1179.
894	Thakurta J., Ripley E.M., and Li, C. (2008) Geochemical constraints on the origin of
895	sulfide mineralization in the Duke Island Complex, southeastern Alaska.
896	Geochemistry, Geophysics, Geosystems 9, 1–34.
897	Tollan P., and Hermann J. (2019). Arc magmas oxidized by water dissociation and
898	hydrogen incorporation in orthopyroxene. Nature Geoscience 12, 667-671.
899	Tomkins A.G., and Evans K.A. (2015) Separate zones of sulfate and sulfide release from
900	subducted mafic oceanic crust. Earth and Planetary Science Letters 428, 73-83.
901	Tomkins A.G., Rebryna K.C., Weinberg R.F., and Schaefer B.F. (2012) Magmatic
902	sulfide formation by reduction of oxidized arc basalt. Journal of Petrology 53, 1537-
903	1567.
904	Villemant B., Jaffrezic H., Joron J.L., and Treuil M. (1981) Distribution coefficients of
905	major and trace-elements-fractional crystallization in the alkali basalt series of
906	Chaine- Des-Puys (Massif Central, France). Geochimica et Cosmochimica Acta 45,
907	1997–2016.
908	Wang C.Y., Wei B., Zhou M., Minh Huu D., and Qi, L. (2018) A synthesis of magmatic
909	Ni-Cu- (PGE) sulfide deposits in the \sim 260 Ma Emeishan large igneous province,
910	SW China and northern Vietnam. Journal of Asian Earth Sciences 154, 162–186.
911	Wang C.Y., Zhou M.F., and Keays, R.R. (2006) Geochemical constraints on the origin of
912	the Permian Baimazhai mafic-ultramafic intrusion, SW China. Contributions to
913	Mineralogy and Petrology152, 309–321.
914	Wang R.M., Liu D.Q., and Yin D.T. (1987) The conditions of controlling metallogeny of
915	Cu-Ni sulfide ore deposits and the orientation of finding ore Hami, Xinjiang, China.
916	Journal of Mineralogy and Petrology 7, 1–152.

- 917 Wang Y.L. (2011) Petrogenesis and mineralization of Heishan intrusion in Beishan area,
- 918 Gansu. M.Sc. thesis, Chang'an University.
- 919 Wei B. (2013) Platinum-group element and Re-Os isotopic compositions of the magmatic
- Ni-Cu sulfide deposits in the Hongqiling-Chajianling-Piaohechuan region, eastern
 part of the Central Asian Orogenic Belt. Ph. D. thesis, University of Chinese
 Academy of Sciences.
- Wei B., Wang C.Y., Arndt N.T., Prichard H.M., and Fisher P.C. (2015) Textural
 relationship of sulfide Ores, PGE, and Sr-Nd-Os isotope compositions of the
 Triassic Piaohechuan Ni-Cu sulfide deposit in NE China. Economic Geology 110,
 2041–2062.
- Wei B., Wang C.Y., Lahaye Y., Xie L., and Cao Y. (2019) S and C Isotope Constraints
 for Mantle-Derived Sulfur Source and Organic Carbon-Induced Sulfide Saturation
 of Magmatic Ni-Cu Sulfide Deposits in the Central Asian Orogenic Belt, North
 China. Economic Geology 114, 787–806.
- Wei B., Wang C.Y., Li C., and Sun Y. (2013) Origin of PGE-depleted Ni-Cu sulfide
 mineralization in the Triassic Hongqiling No. 7 orthopyroxenite intrusion, Central
 Asian Orogenic Belt, northeastern China. Economic Geology 108, 1813–1831.
- Windley B.F., Alexeiev D., Xiao W., Kröner A., and Badarch G. (2007) Tectonic models
 for accretion of the Central Asian Orogenic Belt. Journal of Geological Society 164,
 31–47.
- Windley B.F., Kröner A., Guo J., Qu G., Li Y., and Zhang C. (2002) Neoproterozoic to
 Paleozoic geology of the Altai orogen, NW China: new zircon age data and tectonic
 evolution. Journal of Geology 110, 719–737.
- Wood B.J., Bryndzia L.T., and Johnson K.E. (1990) Mantle oxidation state and its
 relationship to tectonic environment and fluid speciation. Science 248, 337–345.
- Woodland A.B., Kornprobst J., and Tabit A. (2006) Ferric iron in orogenic lherzolite
 massifs and controls of oxygen fugacity in the upper mantle. Lithos 89, 222–241.
- Wu F.Y., Zhao G.C., Sun D.Y., Wilde S.A., and Yang J.H. (2007) The Hulan Group: Its
 role in the evolution of the Central Asian Orogenic Belt of NE China. Journal of
 Asian Earth Sciences 30, 542–556.

- 947 Xia M.-Z., Jiang C.-Y., Li C., and Xia Z.-D. (2013) Characteristics of a newly discovered
- Ni-Cu sulfide deposit hosted in the Poyi ultramafic intrusion, Tarim Craton, NWChina. Economic Geology 108, 1865–1878.
- Xiao W.J., Windley B.F., Badarch G., Sun S., Li J.L., Qin K.Z., and Wang Z.H. (2004a)
 Palaeozoic accretionary and convergent tectonics of the southern Altaids:
 implications for the lateral growth of Central Asia. Journal of Geological Society,
 London 161, 339-342.
- Xiao W.J., Windley B.F., Huang B.C., Han C.M., Yuan C., Chen H.L., Sun M., Sun S.,
 and Li J.L. (2009) End-Permian to mid-Triassic termination of the accretionary
 processes of the southern Altaids: implications for the geodynamic evolution,
 Phanerozoic continental growth, and metallogeny of Central Asia. International
 Journal of Earth Sciences 98, 1189-1287.
- Xiao W.J., Zhang L.C., Qin K.Z., Sun S., and Li J.L. (2004b) Paleozoic accretionary and
 collisional tectonics of the Eastern Tianshan (China): implications for the
 continental growth of central Asia. American Journal of Sciences 304, 370-395.
- Xie W, Song X.Y., Chen L.M., Deng Y.F., Zheng W.Q., Wang Y.S., Ba D.H., Zhang
 X.Q., and Luan Y. (2014) Geochemistry insights on the genesis of the subductionrelated Heishan magmatic Ni-Cu-(PGE) deposit in Gansu, NW China, at the
 southern margin of the Central Asian Orogenic Belt. Economic Geology 109, 1563–
 1583.
- Xie W., Song X., Deng Y., Wang Y., and Ba D. (2012) Geochemistry and petrogenetic
 implications of a Late Devonian mafic- ultramafic intrusion at the southern margin
 of the Central Asian Orogenic Belt. Lithos 144–145, 209–230.
- Xu X., Song S.G., Allen M.B., Ernst R.E., Niu Y.L., and Su L. (2016) An 850–820 Ma
 LIP dismembered during breakup of the Rodinia supercontinent and destroyed by
 Early Paleozoic continental subduction in the northern Tibetan Plateau, NW China.
 Precambrian Research 282, 52–73.
- Xu Y., Chung S., Jahn B., and Wu G. (2001) Petrologic and geochemical constraints on
 the petrogenesis of Permian- Triassic Emeishan flood basalts in southwestern China.
 Lithos 58, 145-168.

Xue S.C., Qin K.Z., Li C., Tang D.M., Mao Y.J., Qi L., and Ripley E.M. (2016)
Geochronological, petrological, and geochemical constraints on Ni-Cu sulfide mineralization in the Poyi ultramafic-troctolitic intrusion in the northeast rim of the
Tarim Craton, Western China. Economic Geology 111, 1465–1484.

- Yang S.H., and Zhou M.F. (2009) Geochemistry of the~ 430-Ma Jingbulake maficultramafic intrusion in Western Xinjiang, NW China: implications for subduction
 related magmatism in the South Tianshan orogenic belt. Lithos 113, 259–273.
- Yang S.H., Zhou M.F., Lightfoot P.C., Malpas J., Qu W.J., Zhou J.B., and Kong D.Y
 (2012) Selective crustal contamination and decoupling of lithophile and chalcophile
 element isotopes in sulfide-bearing mafic intrusions: an example from the

Jingbulake intrusion, Xinjiang, NW China. Chemical Geology 302, 106–118.

- Zhang H.L., Cottrell E., Solheid P.A., Kelley K.A., and Hirschmann M.M. (2018)
 Determination of Fe³⁺/∑Fe of XANES basaltic glass standards by Mössbauer
 spectroscopy and its application to the oxidation state of iron in MORB. Chemical
 Geology 479, 166–175.
- 292 Zhang X., Zhao G., Eizenhöfer P.R., Sun M., Han Y., Hou W., Liu D., Wang B., Liu Q.,
- Xu B., and Yanlin C. (2016) Tectonic transition from Late Carboniferous subduction
 to Early Permian post-collisional extension in the Eastern Tianshan, NW China:
 Insights from geochronology and geochemistry of mafic-intermediate intrusions.
 Lithos 257, 269–281.
- 27 Zhang Z., Li K., Li J., Tang W., Chen Y., and Luo Z. (2015) Geochronology and
 28 geochemistry of the Eastern Erenhot ophiolitic complex: Implications for the
 299 tectonic evolution of the Inner Mongolia-Daxinganling Orogenic Belt. Journal of
 200 Asian Earth Sciences 97, 279–293.
- Zhang Z.C., Mao J.W., Chai F.M., Yan S.H., Chen B.L., and Pirajno F. (2009)
 Geochemistry of the Permian Kalatongke mafic intrusions, northern Xinjiang,
 northwest China: implications for the genesis of magmatic Ni-Cu sulfide deposits.
 Economic Geology 104, 185–203.
- Zhao Y., Xue C., Liu S., Symons D.T.A., Zhao X., Yang Y., and Ke J. (2017) Copper
 isotope fractionation during sulfide-magma differentiation in the Tulaergen
 magmatic Cu deposit, NW China. Lithos 286–287, 206–215.

- Zhao Y., Xue C.J., Zhao X.B., Yang Y.Q., Ke J.J., Zu B., and Zhang G.Z. (2016) Origin
 of anomalously Ni-rich parental magmas and genesis of the Huangshannan Ni-Cu
 sulfide deposit, Central Asian Orogenic Belt, Northwestern China. Ore Geology
 Reviews 77, 57–71.
 Zhou J.B., and Wilde S.A. (2013) The crustal accretion history and tectonic evolution of
- the NE China segment of the Central Asian Orogenic Belt. Gondwana Research 23,
 1365–1377.
- 1015 Zhou M., Lesher C.M., Yang Z., Li J., and Sun M. (2004) Geochemistry and petrogenesis
- 1016 of 270 Ma Ni-Cu-(PGE) sulfide-bearing mafic intrusions in the Huangshan district,
- 1017 Eastern Xinjiang, Northwest China: implications for the tectonic evolution of the
- 1018 Central Asian orogenic belt. Chemical Geology 209, 233-257.
- 1019
- 1020

1021 Figure captions

Fig. 1. (a) The tectonic context of the central Asian orogenic belt (CAOB) relative to other Cratons (modified after Jahn et al., 2000). (b) A simplified geological map of the CAOB (modified after Xiao et al., 2009) showing the mafic-ultramafic intrusions in the CAOB that formed in arc and post-subduction, extensional settings. (c) A geological map of the western segment of the CAOB. (d) A geological map of the eastern segment of the CAOB.

1028

Fig. 2. Chondrite-normalized rare earth element patterns and primitive mantle-normalized trace element patterns for representative mafic-ultramafic intrusions in the CAOB that were emplaced in arc settings. Data sources: Jinbulake (Yang and Zhou, 2009), Heishan (Xie et al., 2012), Erbutu (Peng et al., 2013). Chondrite and primitive mantle values are from Sun and McDonough (1989).

1034

Fig. 3. Chondrite-normalized rare earth element patterns and primitive mantle-normalized trace element patterns for representative mafic-ultramafic intrusions in the CAOB that were emplaced in post-subduction, extensional settings. Data sources: Huangshanxi (Mao et al., 2014), Hongqiling No.2 (Wei, 2013), and Luodong (Su et al., 2011). Chondrite and primitive mantle values are from Sun and McDonough (1989).

1040

Fig. 4. Plot of Mg# versus Cr# (a) and Mg# versus XFe³⁺ (b) for the spinel of the maficultramafic intrusions in the CAOB, and the Dali picrite from the Emeishan large igneous province. Data sources: Jinbulake, Erbutu, Huangshannan and Tulaergen intrusions (this study), Heishan intrusion (Wang, 2011), Baixintan intrusion (this study; Feng et al., 2017), Luodong intrusion (Su et al., 2011), Hongqiling No.1 and No.2 intrusions (Cao et al., 2019; Wei et al., 2019), Dali picrite (Kamenetsky et al., 2012; Liu et al., 2017).

1047

Fig. 5. Histogram of δ^{34} S values of sulfides from the Jinbulake, Erbutu, Baixintan and Tulaergen intrusions in the CAOB. The δ^{34} S values of MORB-type mantle are from Labidi et al. (2014).

1051

Fig. 6. Variation of V/Sc of the primary magma against the degrees of partial melting (F) at given fO_2 (Lee et al., 2005). It is assumed that the mafic-ultramafic intrusions in the CAOB were derived from magmas produced by ~15 to ~20% of partial melting (indicated by the grey shaded area) of the mantle wedge in the spinel stability field.

Fig. 7. Plot of XCr^{3+} of spinel versus $\ln Kd_{Mg/Fe}^{Ol-Spl}$ for the mafic-ultramatic intrusions in the CAOB, and the Dali picrite in the Emeishan large igneous province. Data sources are the same as those in Fig. 4.

1060

Fig. 8. (a) Comparison of the estimated magma fO_2 of the mafic-ultramatic intrusions in 1061 1062 the CAOB and the Dali picrite in the Emeishan large igneous province with the fO_2 of MORBs (FMQ-1 to FMQ+0.5) and arc basalts (FMQ+0.5 to FMQ+6). Data sources: 1063 MORBs (Cottrell and Kelley, 2011; Zhang et al., 2018), arc basalts (Woodland et al., 1064 2006). (b) Plot of the magma fO_2 versus the Fo contents of olivine for the mafic-1065 ultramafic intrusions in the CAOB and the Dali picrite in the Emeishan large igneous 1066 1067 province. The error bar in (b) represents the uncertainty (FMQ ± 0.4) of calculated magma fO_2 based on the olivine-spinel oxygen barometer (*c.f.*, Ballhaus et al., 1991). The dashed 1068 1069 line outlines the data for the intrusions with tholeiitic, parental magmas. 1070 Fig. 9. (a) Comparison of the magma fO_2 calculated based on olivine-spinel oxygen 1071

Fig. 9. (a) Comparison of the magma fO_2 calculated based on olivine-spinel oxygen barometer and the partitioning of V in olivine showing the good agreement of the results obtained by two different methods. The error bars represent the uncertainty of magma fO_2 calculated based on the two methods. (b) Plot of the magma fO_2 calculated based on the olivine-spinel oxygen barometer versus the V/Sc of olivine. There is an overall negative relationship between the magma fO_2 and the V/Sc of olivine. The error bar represents 1 σ standard deviation of the measured V/Sc of olivine.

1078

Fig. 10. Plot of Alz (percentage of tetrahedral sites occupied by Al) versus wt.% TiO_2 of clinopyroxene from the mafic-ultramafic intrusions in the CAOB and the Emeishan large

1081 igneous province. The trends of the arc and rift cumulate are modified after Loucks1082 (1990).

1083

1084

Fig. 11. Comparison of δ^{34} S values of sulfides and magma fO_2 among the mafic-1085 ultramafic intrusions in the CAOB, the Jinchuan Ni-Cu sulfide deposits in the southern 1086 margin of the North China Craton, and the Ni-Cu sulfide deposits in the Emeishan large 1087 1088 igneous province. The mafic-ultramafic intrusions in the CAOB overall have fO_2 >FMQ+1 and $\delta^{34}S$ similar to the MORB mantle value (-1.6 to +0.6%; Labidi et al., 1089 2013, 2014), whereas the Ni-Cu sulfide deposits in the intraplate settings have relatively 1090 low fO_2 and high $\delta^{34}S$ of sulfides. Data sources: Jinbulake, Erbutu, Baixintan and 1091 Tulaergen intrusions (this study), Heishan intrusion (Xie et al., 2014), Honggiling No.7 1092 intrusion (Wei et al., 2019), Luodong intrusion (Su et al., 2015), Povi intrusion (Xia et al., 1093 1094 2013), Huangshannan intrusion (Zhao et al., 2016), Huangshandong and Huangshanxi 1095 intrusions (Wang et al., 1987), Jinchuan intrusion (Ripley et al., 2005; Duan et al., 2016), 1096 the intrusions in the Emeishan large igneous province (Wang et al., 2018).

1097

Fig. 12. Comparison of magma fO_2 values calculated based on the olivine-spinel oxygen barometer with those calculated based on the Fe-Ni exchange between olivine and sulfide melt for the Baixintan (BXT), Huangshannan (HSN), Huangshandong (HSD), Huangshanxi (HSX), Tulaergen (TLEG), and Hongqiling No.1 (HQL) intrusions in the CAOB. The values based on the Fe-Ni exchange between olivine and sulfide liquids are much lower than those based on the olivine-spinel oxygen barometer.

	a 1 1	0.101	$\delta^{34}S\%$		a 1 3 4	0.101	$\delta^{34}S\%$
Analysis No.	Sample No.	Sulfides	(V-CDT)	Analysis No.	Sample No.	Sulfides	(V-CDT)
Jinbulake intrusion				15	18EBT-11	pentlandite	6.2
1	QB-13	pyrrhotite	0.6	16	18EBT-11	pyrrhotite	6.2
2	QB-13	pyrrhotite	0.3	17	18EBT-11	pyrrhotite	6.4
3	QB-13	pyrrhotite	0.0	18	18EBT-5	pentlandite	5.5
4	QB-13	pyrrhotite	0.5	19	18EBT-5	pentlandite	5.5
5	QB-13	pyrrhotite	0.9	20	18EBT-5	pyrrhotite	5.8
6	QB2-102	pyrrhotite	0.9	21	18EBT-5	pyrrhotite	5.3
7	QB2-102	pyrrhotite	0.8	22	18EBT-5	pyrrhotite	5.6
8	QB2-102	pyrrhotite	0.8	23	18EBT-5	chalcopyrite	6.0
9	QB2-102	pyrrhotite	1.3	24	18EBT-5	pentlandite	6.0
10	QB2-78	pyrrhotite	0.1	25	18EBT-5	pentlandite	5.6
11	QB2-78	pyrrhotite	0.2	Baixintan intru	ision		
12	QB2-78	pyrrhotite	0.7	1	19BXT-4	chalcopyrite	0.4
13	QB2-78	pyrrhotite	0.5	2	19BXT-4	chalcopyrite	0.3
14	QB-43	pyrrhotite	0.3	3	19BXT-4	pyrrhotite	-0.1
15	QB-43	pentlandite	0.3	4	19BXT-4	pyrrhotite	0.5
16	QB-43	pentlandite	0.3	5	19BXT-4	chalcopyrite	1.1
17	QB-43	pyrrhotite	0.6	6	19BXT-6	pyrite	0.7
18	QB-43	pyrrhotite	0.7	7	19BXT-6	pyrite	0.5
19	QB-43	pentlandite	0.6	8	19BXT-6	pyrite	0.6
20	QB-43	pyrrhotite	0.7	9	19BXT-14	chalcopyrite	0.6
21	QB-65	chalcopyrite	0.9	10	19BXT-14	chalcopyrite	0.1
22	QB-65	chalcopyrite	0.9	11	19BXT-14	pyrrhotite	-0.7
23	QB-65	pyrrhotite	0.6	12	19BXT-14	pyrrhotite	0.1
24	QB-65	pyrrhotite	0.4	13	19BXT-ZK-15	chalcopyrite	1.2
25	QB-65	chalcopyrite	1.3	14	19BXT-ZK-15	pyrrhotite	0.0
26	QB-65	pentlandite	0.7	15	19BXT-ZK-15	chalcopyrite	-0.4
27	QB-65	pyrrhotite	0.9	16	19BXT-ZK-15	chalcopyrite	0.2
28	QB-65	pyrrhotite	0.6	17	19BXT-ZK-15	chalcopyrite	-0.1
Erbutu intrusic	on			18	19BXT-ZK-15	pentlandite	-0.3
1	18EBT-10	pyrrhotite	7.5	19	19BXT-ZK-15	pentlandite	-0.4
2	18EBT-10	pyrrhotite	7.4	Tulaergen intro	usion		
3	18EBT-10	chalcopyrite	7.3	1	TLEG-16	pyrrhotite	0.5
4	18EBT-10	chalcopyrite	5.9	2	TLEG-16	pyrrhotite	0.9
5	18EBT-11	chalcopyrite	6.7	3	TLEG-16	pyrrhotite	0.3
6	18EBT-11	pentlandite	6.2	4	TLEG-16	pentlandite	0.4
7	18EBT-11	pentlandite	6.0	5	TLEG-16	pyrrhotite	0.3
8	18EBT-11	chalcopyrite	6.7	6	TLEG-19	pyrrhotite	0.2
9	18EBT-11	chalcopyrite	6.9	7	TLEG-19	pyrrhotite	-0.2
10	18EBT-11	pentlandite	6.2	8	TLEG-19	pyrrhotite	0.3
11	18EBT-11	pentlandite	6.7	9	TLEG-19	pyrrhotite	0.4
12	18EBT-11	chalcopyrite	6.9	10	TLEG-26	chalcopyrite	0.5
13	18EBT-11	pentlandite	6.4	11	TLEG-26	pyrrhotite	0.2
14	18EBT-11	pentlandite	5.9				

Table 1 S isotopic compositions of the sulfides in the rocks from the Jinbulake, Erbutu, Baixintan and Tulaergen intrusions in the central Asian orogenic belt



This is the peer-reviewed, final accepted version for American Mineralogist, published by the Mineralogical Society of America. The published version is subject to change. Cite as Authors (Year) Title. American Mineralogist, in press. DOI: https://doi.org/10.2138/am-2020-7351. http://www.minsocam.org/



Always consult and cite the final, published document. See http://www.minsocam.org or GeoscienceWorld

This is the peer-reviewed, final accepted version for American Mineralogist, published by the Mineralogical Society of America. The published version is subject to change. Cite as Authors (Year) Title. American Mineralogist, in press. DOI: https://doi.org/10.2138/am-2020-7351. http://www.minsocam.org/



Always consult and cite the final, published document. See http://www.minsocam.org or GeoscienceWorld











Fig. 8





Fig. 11



