

**ACTINIDES IN GEOLOGY, ENERGY, AND THE ENVIRONMENT**

**Uranium scavenging during mineral replacement reactions†**

**KAN LI<sup>1</sup>, ALLAN PRING<sup>2,4</sup>, BARBARA ETSCHMANN<sup>1,2,7</sup>, EDELTRAUD MACMILLAN<sup>3</sup>, YUNG NGOTHAI<sup>1</sup>,  
BRIAN O'NEILL<sup>1</sup>, ANTHONY HOOKER<sup>1,5</sup>, FRED MOSSELMANS<sup>6</sup> AND JOËL BRUGGER<sup>2,7,\*</sup>**

<sup>1</sup>School of Chemical Engineering, The University of Adelaide, 5000, South Australia, Australia

<sup>2</sup>Division of Mineralogy, South Australian Museum, North Terrace, 5000, Adelaide, South Australia, Australia

<sup>3</sup>School of Earth and Environmental Sciences, The University of Adelaide, 5000, South Australia, Australia

<sup>4</sup>School of Chemical and Physical Sciences, Flinders University, 5001, South Australia, Australia

<sup>5</sup>Radiation Health, Radiation Protection Branch, Environment Protection Authority, Victoria SQ, 5000, Adelaide, South Australia, Australia

<sup>6</sup>Diamond Light Source Ltd., Diamond House, Harwell Science and Innovation Campus, Didcot, Oxfordshire OX11 0DE, U.K.

<sup>7</sup>School of Earth, Atmosphere, and Environment, Monash University, 3800, Clayton, Victoria, Australia

**ABSTRACT**

Interface coupled dissolution-reprecipitation reactions (ICDR) are a common feature of fluid-rock interaction during crustal fluid flow. We tested the hypothesis that ICDR reactions can play a key role in scavenging minor elements by exploring the fate of U during the experimental sulfidation of hematite to chalcopyrite under hydrothermal conditions (220–300 °C). The experiments where U was added, either as solid  $\text{UO}_{2+x}(\text{s})$  or as a soluble uranyl complex, differed from the U-free experiments in that pyrite precipitated initially, before the onset of chalcopyrite precipitation. In addition, in  $\text{UO}_{2+x}(\text{s})$ -bearing experiments, enhanced hematite dissolution led to increased porosity and precipitation of pyrite+magnetite within the hematite core, whereas in uranyl nitrate-bearing experiments, abundant pyrite formed initially, before being replaced by chalcopyrite. Uranium scavenging was mainly associated with the early reaction stage (pyrite precipitation), resulting in a thin U-rich line marking the original hematite grain surface. This “line” consists of nanocrystals of  $\text{UO}_{2+x}(\text{s})$ , based on chemical mapping and XANES spectroscopy. This study shows that the presence of minor components can affect the pathway of ICDR reactions. Reactions between U- and Cu-bearing fluids and hematite can explain the Cu-U association prominent in some iron oxide-copper-gold (IOCG) deposits.

**Keywords:** Uranium, scavenging, IOCG deposits, experiment, sulfidation reaction, interface coupled dissolution-reprecipitation reactions