

## Chemistry-dependent Raman spectral features of glauconite and nontronite: Implications for mineral identification and provenance analysis

RAPHAEL J. BAUMGARTNER<sup>1,2,\*</sup>, JAVIER CUADROS<sup>3</sup>, JOSEPH MICHALSKI<sup>4</sup>, BOBBY PEJČIĆ<sup>1</sup>,  
CARSTEN LAUKAMP<sup>1</sup>, SIYU HU<sup>1,†</sup>, AND JULIEN BOURDET<sup>5</sup>

<sup>1</sup>CSIRO Mineral Resources, Australian Resources Research Centre, Kensington, Western Australia 6151 Australia

<sup>2</sup>School of Biological, Earth and Environmental Sciences, The University of New South Wales, Kensington, New South Wales 2033, Australia

<sup>3</sup>Department of Earth Sciences, The Natural History Museum, London SW7 5BD, U.K.

<sup>4</sup>Department of Earth Sciences and Laboratory for Space Research, University of Hong Kong, Hong Kong, China

<sup>5</sup>CSIRO Energy, Australian Resources Research Centre, Kensington, Western Australia 6151, Australia

### ABSTRACT

This study provides a comprehensive Raman spectral characterization of nontronite and glauconite-nontronite mixed-layer phases from seafloor hydrothermal fields. These 2:1 phyllosilicates, which show isomorphous cation exchange between  $\text{Mg}^{2+}+\text{Fe}^{2+}$  and  $\text{Fe}^{3+}+\text{Al}^{3+}$  in the dioctahedral sheets, exhibit three diagnostic Raman peaks in the low wavenumber region ( $\nu_1 \sim 241\text{--}257\text{ cm}^{-1}$ ;  $\nu_2 \sim 600\text{--}606\text{ cm}^{-1}$ ;  $\nu_3 \sim 690\text{ cm}^{-1}$ ), and one peak at  $\sim 3548\text{--}3570\text{ cm}^{-1}$  ( $\nu_4$ ). With increasing  $(\text{Mg}^{2+}+\text{Fe}^{2+})_{\text{oct}}$ , the presumed stretching band of octahedral OH-O bonds ( $\nu_1$ ) is displaced to a higher wavenumber, whereas the stretching band of tetrahedral Si-O-Si bonds ( $\nu_2$ ) is shifted to a lower wavenumber. Peak  $\nu_4$ , which relates to O-H bonds of hydroxyls linked to octahedral cations, shows a downshift with increasing  $(\text{Mg}^{2+}+\text{Fe}^{2+})_{\text{oct}}$ . The band  $\nu_4$  can be mathematically fitted by three bands, two of which strongly correlate with the cation occupancy in the octahedral sheets; i.e., vibrations of hydroxyls linked to trivalent cations ( $\text{Fe}^{3+}$  and  $\text{Al}^{3+}$ ) are mainly represented by a band at  $\sim 3560\text{--}3573\text{ cm}^{-1}$ , whereas divalent cations ( $\text{Mg}^{2+}$  and  $\text{Fe}^{2+}$ ) mainly contribute to a band at  $\sim 3538\text{--}3540\text{ cm}^{-1}$ . This result is consistent with theoretical considerations for dioctahedral phyllosilicates, which predict for the incorporation of  $\text{Mg}^{2+}$  and  $\text{Fe}^{2+}$  a weakening/lengthening of O-H bonds in the OH groups, accounting for a downshift of the O-H vibrations. Hence, this is one of the first studies that trace how even subtle chemical modifications in phyllosilicates influence Raman spectral features. The reported findings have implications for mineral identification and provenance analysis, such as during surface exploration on Mars, where compositionally diverse phyllosilicates occur.

**Keywords:** Phyllosilicates, glauconite, nontronite, peak shift