

Lakebogaite, $\text{CaNaFe}_2^3\text{H}(\text{UO}_2)_2(\text{PO}_4)_4(\text{OH})_2(\text{H}_2\text{O})_8$, a new uranyl phosphate with a unique crystal structure from Victoria, Australia

STUART J. MILLS,^{1,2,3,*} WILLIAM D. BIRCH,² UWE KOLITSCH,⁴ W. GUS MUMME,³ AND IAN E. GREY³

¹School of Earth Sciences, University of Melbourne, Parkville, Victoria, 3010, Australia

²Geosciences, Museum Victoria, GPO Box 666, Melbourne, Victoria, 3001, Australia

³CSIRO Minerals, Box 312, Clayton South, Victoria, 3169, Australia

⁴Mineralogisch-Petrographische Abt., Naturhistorisches Museum Wien, Burgring 7, A-1010 Wien, Austria

ABSTRACT

Lakebogaite, ideally $\text{CaNaFe}_2^3\text{H}(\text{UO}_2)_2(\text{PO}_4)_4(\text{OH})_2(\text{H}_2\text{O})_8$, is a new Ca–Na–Fe uranyl phosphate mineral from a quarry in Upper Devonian granite near Lake Boga, northern Victoria, Australia. It is associated with Na-analogue of meurigite (IMA 2007-024), torbernite, and saléeite on a matrix of microcline, albite, smoky quartz, and muscovite. Lakebogaite occurs as bright lemon-yellow transparent prismatic crystals up to 0.4 mm across. The crystals have a vitreous luster and a pale yellow streak. Mohs hardness is about 3. The fracture is even to conchoidal. In transmitted light, the mineral is pale yellow with very weak pleochroism: X = yellow, Y = grayish yellow, Z = grayish yellow: dispersion $r > v$, strong. Lakebogaite crystals are biaxial (+), with slightly variable refractive indices within the ranges: $n_\alpha = 1.650(2)$ – $1.652(2)$, $n_\beta = 1.660(4)$ – $1.664(3)$, $n_\gamma = 1.681(3)$ – $1.686(2)$, measured using white light, and with $2V_{\text{meas}} = 80$ – 85° and $2V_{\text{calc}} = 70$ – 74° . Orientation: Y = **b**; crystals are elongated along [010], resulting in straight extinction. The empirical chemical formula (mean of nine electron microprobe analyses) calculated on the basis of 30 anions is $(\text{Ca}_{1.00}\text{Na}_{0.80}\text{Sr}_{0.10})_{\Sigma 1.90}(\text{Fe}_{1.85}^{3+}\text{Al}_{0.30})_{\Sigma 2.15}(\text{UO}_2)_{1.80}(\text{PO}_4)_{4.07}(\text{OH},\text{H}_2\text{O})_{10.12}$. Lakebogaite is monoclinic, space group *Cc*, $a = 19.6441(5)$, $b = 7.0958(2)$, $c = 18.7029(5)$ Å, $\beta = 115.692(1)^\circ$, $V = 2349.3(7)$ Å³, $Z = 4$. The seven strongest reflections in the powder X-ray diffraction pattern are [d_{obs} in Å (*hkl*): 6.60 (100) (110), 3.16 (40) (514, 604), 4.07 (20) (404), 3.80 (20) (314), 3.56 (20) (020, 312), 3.31 (20) (114, 220), 2.797 (20) (006)]. The crystal structure was solved from single-crystal X-ray diffraction data and refined to $R_1 = 0.038$ on the basis of 5222 unique reflections with $F > 4\sigma F$. It comprises pairs of edge-shared UO_7 pentagonal bipyramids that are inter-linked via corner-sharing with PO_4 tetrahedra, to form chains parallel to the **c**-axis. Each UO_7 polyhedron also shares one of its edges with another PO_4 tetrahedron. The $(\text{UO}_2)_2(\text{PO}_4)_4$ chains are cross-linked via corner-sharing between the PO_4 tetrahedra and $\text{Fe}^{3+}\text{O}_4(\text{OH})_2$ octahedra. The octahedra join together by corner-sharing via OH anions to form chains parallel to **b**. The Na^+ and Ca^{2+} cations, and 4 water molecules occupy eight-sided channels along [010]. The remaining water molecules occupy large ten-sided channels directed along [001] and intersecting the [010] channels. The mineral is named for the nearest township.

Keywords: Lakebogaite, new mineral, uranyl phosphate, Lake Boga quarry, north-western Victoria, Australia, crystal structure, $(\text{UO}_2)_2(\text{PO}_4)_4$ chains