

## **Multiple titanium substitutions in biotites from high-grade metapelitic xenoliths (Euganean Hills, Italy): Complete crystal chemistry and appraisal of petrologic control**

**RAFFAELE SASSI,<sup>1,\*</sup> GIUSEPPE CRUCIANI,<sup>2</sup> CLAUDIO MAZZOLI,<sup>1,3</sup> LUCA NODARI,<sup>4</sup> AND JOHN CRAVEN<sup>5</sup>**

<sup>1</sup>Dipartimento di Geoscienze, Università di Padova, Via Giotto 1, I-35121 Padova, Italy

<sup>2</sup>Dipartimento di Scienze della Terra, Università di Ferrara, Via Saragat 1, I-44100 Ferrara, Italy

<sup>3</sup>Istituto di Geoscienze e Georisorse, CNR, C.so Garibaldi 37, I-35121 Padova, Italy

<sup>4</sup>Dipartimento di Scienze Chimiche, Università di Padova, Via Marzolo 1, I-35131 Padova, Italy

<sup>5</sup>School of Geosciences, Grant Institute, The King's Buildings, West Mains Road, Edinburgh EH9 3JW, U.K.

### **ABSTRACT**

Biotites from metapelitic xenoliths included within trachytes from the Euganean Hills (Italy) were analyzed by single-crystal X-ray diffraction (XRD), electron microprobe (EMP), scanning electron microscope (SEM), secondary ion mass spectrometry (SIMS), and Mössbauer spectroscopy. These biotites are Ti-rich and occur in gneissic xenoliths that underwent regional high-*T*/low-*P* metamorphism, at about 750 °C, followed by pyrometamorphism during incorporation in the melt at temperatures close to 950 °C.

Biotites are zoned, with TiO<sub>2</sub> content ranging from 6.79 (cores) to 8.14 wt% (rims). SIMS measurements show that the H<sub>2</sub>O content is in the range 2.88–4.08 wt%. The simultaneous occurrence of high-Ti and high-H<sub>2</sub>O contents, and the main cation substitutions based on EMP analyses suggest that the role of Ti-oxy in these biotites is less important than Ti-vacancy and Ti-Tschermak substitutions. Single-crystal XRD confirms that the Ti-oxy exchange was indeed effective but not the dominant substitution mechanism.

Based on our data and those taken from literature on petrologically well-constrained systems, we propose that there is a petrologic control on the type of Ti-substitution mechanisms. We consider two types of petrologic groupings for biotites: (1) group A consisting of biotites from H<sub>2</sub>O-free or H<sub>2</sub>O-poor petrologic environments (e.g., volcanic rocks, ultrabasic xenoliths, and crustal xenoliths in which biotite underwent incongruent melting): Ti substitution in these biotites occurs via Ti-oxy predominantly, or more specifically Fe<sup>3+</sup>-Ti-oxy; and (2) group B consisting of biotites from H<sub>2</sub>O-rich petrologic environments (e.g., metamorphic rocks and crustal granitoids): Ti-vacancy, or more specifically Fe<sup>3+</sup>-Ti-vacancy, is the dominant mechanism in them.

It is concluded that during high-grade metamorphism the dominating type of Ti substitution in biotite is controlled by H<sub>2</sub>O activity.

**Keywords:** Biotite, Ti-oxy substitution, SCXRD, SIMS