

LETTER

New high-pressure B2 phase of FeS above 180 GPa

**NAGAYOSHI SATA,^{1,*} HIROAKI OHFUJI,² KEI HIROSE,^{1,3} HISAO KOBAYASHI,⁴ YASUO OHISHI,⁵ AND
NAOHISA HIRAO⁵**

¹Institute for Research on Earth Evolution, Japan Agency for Marine-Earth Science and Technology, 2-15 Natsushima-cho, Yokosuka, Kanagawa
237-0061, Japan

²Geodynamics Research Center, Ehime University, 2-5 Bunkyo-cho, Matsuyama, Ehime 790-8577, Japan

³Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro, Tokyo 152-8551, Japan

⁴Graduate School of Material Science, University of Hyogo, 3-2-1 Kouto, Kamigori-cho, Ako-gun, Hyogo 678-1297, Japan

⁵Japan Synchrotron Radiation Research Institute, 1-1-1 Kouto, Sayo-cho, Sayo-gun, Hyogo 679-5198, Japan

ABSTRACT

FeS exhibits extensive polymorphism at high pressure and temperature. All with NiAs-type (B8) or closely related structures. Here we report a new phase transition from FeS VI to CsCl-type (B2) phase (FeS VII) above 180 GPa based on the synchrotron X-ray diffraction (XRD) measurements. A significant volume reduction by 3.0% was observed at the phase transition, due to an increase in the coordination number from six to eight. Present results suggest that a substantial amount of sulfur may be incorporated into an Fe-Ni alloy with *bcc* structure in the Earth's inner core.

Keywords: Earth's core, troilite, high pressure, phase transformation, iron sulfide