

## **Low-pressure clino- to high-pressure clinoenstatite phase transition: A phonon-related mechanism**

**YONGGANG G. YU<sup>1,\*</sup> AND RENATA M. WENTZCOVITCH<sup>2,†</sup>**

<sup>1</sup>Department of Chemistry, Minnesota Supercomputing Institute, University of Minnesota, Minneapolis, Minnesota 55455, U.S.A.

<sup>2</sup>Department of Chemical Engineering and Materials Sciences, Minnesota Supercomputing Institute, University of Minnesota, Minneapolis, Minnesota 55455, U.S.A.

### **ABSTRACT**

We have investigated by first principles the compressional behavior of low-pressure (LP) and high-pressure (HP) MgSiO<sub>3</sub> clinoenstatite. We have carefully examined unit-cell shapes, chain angles, and polyhedral volume responses, such as angle variances and quasi-elongations, under pressure at room temperature. We have observed opposite behavior of the tetrahedra in the S-rotated and O-rotated chains with pressure in the LP phase, with a slight increase (decrease) in angle variance and quasi-elongation in the former (latter). Inspection of zone center modes of both phases under pressure reveals a transition path that converts the S-rotated chain in the LP phase into the O-rotated chain in the HP phase. This conversion is related to a slight softening of an  $A_g$  “metastable” Raman mode under pressure.

**Keywords:** Phase transition mechanism, low-pressure clinoenstatite, high-pressure clinoenstatite, Raman, IR, phonon frequency